Completeness of the propositions-as-types interpretation of intuitionistic logic into illative combinatory logic[1]

Wil Dekkers*
Martin Bunder†
Henk Barendregt‡

October 1995

Abstract

Illative combinatory logic consists of the theory of combinators or lambda calculus extended by extra constants (and corresponding axioms and rules) intended to capture inference. In a preceding paper, BBD[1993], we considered 4 systems of illative combinatory logic that are sound for first order intuitionistic propositional and predicate logic. The interpretation from ordinary logic into the illative systems can be done in two ways: following the propositions-as-types paradigm, in which derivations become combinators, or in a more direct way, in which derivations are not translated. Both translations are closely related in a canonical way. In the cited paper we proved completeness of the two direct translations. In the present paper we prove that also the two indirect translations are complete. Moreover we shall prove that one of the systems is also complete for predicate calculus with higher type functions.

[1]Research supported by the Australian Research Council.

*Faculty of Mathematics and Computer Science, Catholic University, Nijmegen, The Netherlands, e-mail: wil@cs.kun.nl.

†Faculty of Informatics, Department of Mathematics, University of Wollongong, NSW Australia, e-mail: m.bunder@uow.edu.au.

‡Faculty of Mathematics and Computer Science, Catholic University, Nijmegen, The Netherlands, e-mail: henk@cs.kun.nl.

1. Introduction

This paper is a continuation of a preceding paper, BBD[1993], by the same authors. We will refer to that paper mostly as B; so for example Proposition B3.10 means Proposition 3.10 in BBD[1993]. As the present paper is a continuation of that paper, we will not repeat all the definitions and results here but will give only a very short summary.

In Definition B2.3 we defined 4 systems of illative combinatory logic $\mathcal{I}\mathbf{P}, \mathcal{I}\mathbf{\Xi}, \mathcal{I}\mathbf{F}$ and $\mathcal{I}\mathbf{G}$. We interpreted intuitionistic propositional (the $\{\rightarrow\}-$fragment) and predicate logic (the $\{\rightarrow\forall\}-$fragment) into illative systems in two ways: following the propositions-as-types paradigm, in which derivations become combinators, and in a more direct way, in which derivations are not translated.

We had as aim to prove

(i) $\Delta \vdash_{\mathrm{PROP}} \varphi \Leftrightarrow \forall r[\ldots_1, [\Delta]^1_r \vdash_{\mathcal{I}\mathrm{P}} [\varphi]^1_r]$;
(ii) $\Delta \vdash_{\mathrm{PRED}} \varphi \Leftrightarrow \forall r[\ldots_1, [\Delta]^1_r \vdash_{\mathcal{I}\Xi} [\varphi]^1_r]$;
(iii) $\Delta \vdash_{\mathrm{PROP}} \varphi \Leftrightarrow \forall r \exists M[\ldots_2, [\Delta]^2_r \vdash_{\mathcal{I}\mathrm{F}} [\varphi]^2_r M]$;
(iv) $\Delta \vdash_{\mathrm{PRED}} \varphi \Leftrightarrow \forall r \exists M[\ldots_2, [\Delta]^2_r \vdash_{\mathcal{I}\mathrm{G}} [\varphi]^2_r M]$;

where $\ldots_i$ stands for $\Gamma^i_r(\Delta, \varphi)$ in the cases of PROP and for $\Gamma^{i,+}_{r,s}, \Gamma(\Delta, \varphi)$ in the cases of PRED.

The proofs of soundness ($\Rightarrow$)were easy. (cf. Propositions B2.11 and B2.14). In Propositions B3.14 and B3.11 we proved completeness for the 2 direct translations $[\ \ ]^1 : \mathrm{PROP} \rightarrow \mathcal{I}\mathbf{P}$ and $[\ \ ]^1 : \mathrm{PRED} \rightarrow \mathcal{I}\mathbf{\Xi}$, ((i)$\Leftarrow$ and (ii)$\Leftarrow$), by specialising to $r = \mathbf{I}$. In the present paper we prove completeness for the 2 indirect translations $[\ ]^2 : \mathrm{PROP} \rightarrow \mathcal{I}\mathbf{F}$ and $[\ ]^2 : \mathrm{PRED} \rightarrow \mathcal{I}\mathbf{G}$, ((iii)$\Leftarrow$ and (iv)$\Leftarrow$), also by specialising to $r = \mathbf{I}$. In section 5 we point out that completeness for all 4 cases can be proved also by specialising to $r = \mathbf{K}$.

The completeness proof for the direct translation $[\ ]^1 : \mathrm{PRED} \rightarrow \mathcal{I}\mathbf{\Xi}$ ((ii)$\Leftarrow$) was given in the following way. We had to show

$$\Gamma^{1,+}_s, [\Delta]^1, \Gamma(\Delta, \varphi) \vdash_{\mathcal{I}\Xi} [\varphi]^1 \ \Rightarrow\ \Delta \vdash_{\mathrm{PRED}} \varphi.$$

(c.f. page B781.) Write $\Gamma' = \Gamma^{1,+}_s, [\Delta]^1, \Gamma(\Delta, \varphi)$. First we defined in B3.2 a set $\mathcal{O}$ in order to analyze the terms M such that $\Gamma' \vdash_{\mathcal{I}\Xi} M$. We defined $\overline{\mathcal{O}} = \{M \mid \exists N \in \mathcal{O} \mid N =_{\beta\eta} M\}$ and it was clear that $\Gamma' \subseteq \overline{\mathcal{O}}$. We showed in Proposition B3.10 that $\mathcal{O}$ is an invariant (modulo $\beta\eta$-equality) for derivations in $\mathcal{I}\mathbf{\Xi}$:

$$\Gamma \vdash_{\mathcal{I}\Xi} M, \Gamma \subseteq \overline{\mathcal{O}} \Rightarrow M \in \overline{\mathcal{O}}.$$

Using this result we then proved in Proposition B3.11

$$\Gamma' \vdash_{\mathcal{I}\Xi} [\varphi]^1 \ \Rightarrow\ \Delta \vdash_{\mathrm{PRED}} \varphi.$$

The completeness proof for $[\ ]^1 : \mathrm{PROP} \rightarrow \mathcal{I}\mathbf{P}$ ((i)$\Leftarrow$) was done in a similar but easier way.

In the present paper we will prove completeness of $[\ ]^2 : \mathrm{PROP} \to \mathcal{I}\mathbf{F}$ and $[\ ]^2 : \mathrm{PRED} \to \mathcal{I}\mathbf{G}$ ((iii)$\Leftarrow$ and (iv)$\Leftarrow$).

We start with the most difficult case, the proof of completeness of $[\ ]^2 : \mathrm{PRED} \to \mathcal{I}\mathbf{G}$. We define in Definition 2.2 a set, denoted by $\mathcal{O}_G$ to distinguish it from $\mathcal{O}$, and we will show in Proposition 2.10 that $\mathcal{O}_G$ is an invariant (modulo $\beta\eta$-equality) for derivations in $\mathcal{I}\mathbf{G}$:

$$\Gamma \vdash_{\mathcal{I}\mathrm{G}} M, \Gamma \subseteq \overline{\mathcal{O}_G} \ \Rightarrow \ M \in \overline{\mathcal{O}_G}.$$

The invariant $\mathcal{O}_G$ is more complicated than $\mathcal{O}$ and we cannot prove completeness directly using this invariant, as in B3.11 for $\mathcal{I}\mathbf{\Xi}$ relative to PRED. But we shall use $\mathcal{O}_G$ in order to show in Lemma 2.18 that in a proof of $\Gamma \vdash_{\mathcal{I}\mathrm{G}} M$, where $\Gamma \subseteq \overline{\mathcal{O}}$, all cut formulas can be eliminated. Then we define in Definition 2.21 a second invariant $\mathcal{O}_1$, similar to $\mathcal{O}$ in B3.2 and using this invariant we prove in 2.22 - 2.25 the completeness in a way similar to the proof of completeness of $[\ ]^1 : \mathrm{PRED} \to \mathcal{I}\mathbf{\Xi}$ in B3.2 - B3.11.

Completenes of $\mathcal{I}\mathbf{F}$ for PROP is easier. There is no need for cut elimination and we give the proof directly, using an invariant $\mathcal{O}_2$ similar to $\mathcal{O}$ and $\mathcal{O}_1$.

In the last part of the paper we shall introduce another logic system: Predicate Calculus with Higher Type Functions, notation PRED$\lambda\!\to$. We shall prove in Proposition 4.12 that $\mathcal{I}\mathbf{G}$ is also complete for PRED$\lambda\!\to$ under the second translation $[\ ]^2$.

2. Completeness of $\mathcal{I}\mathbf{G}$ for PRED

We will show for $\varphi \in \mathbb{F}_{\mathrm{PRED}},\ \Delta \subseteq \mathbb{F}_{\mathrm{PRED}}$:

$$\forall r[\ \Gamma^{2,+}_{r,s}, [\Delta]^2_r, \Gamma(\Delta, \varphi) \vdash_{\mathcal{I}\mathrm{G}} [\varphi]^2_r M \text{ some } M\ ] \ \Rightarrow \ \Delta \vdash_{\mathrm{PRED}} \varphi.$$

It is sufficient to show

$$\Gamma^{2,+}_{\mathsf{I},s}, [\Delta]^2_{\mathsf{I}}, \Gamma(\Delta, \varphi) \vdash_{\mathcal{I}\mathrm{G}} [\varphi]^2_{\mathsf{I}} M \ \Rightarrow \ \Delta \vdash_{\mathrm{PRED}} \varphi,$$

We omit the index I in the notation and we prove

$$\Gamma^{2,+}_{s}, [\Delta]^2, \Gamma(\Delta, \varphi) \vdash_{\mathcal{I}\mathrm{G}} [\varphi]^2 M \ \Rightarrow \ \Delta \vdash_{\mathrm{PRED}} \varphi.$$

The proof is again in several steps. First we define in 2.2 an *invariant* $\mathcal{O}_G$ in order to analyze the terms N such that $\Gamma^{2,+}_s, [\Delta]^2, \Gamma(\Delta, \varphi) \vdash_{\mathcal{I}\mathrm{G}} N$. Using this invariant we show in Lemma 2.18 that in a proof of $\Gamma^{2,+}_s, [\Delta]^2, \Gamma(\Delta, \varphi) \vdash_{\mathcal{I}\mathrm{G}} N$ all cuts can be eliminated . Then we define in Definition 2.21 a second invariant $\mathcal{O}_1$, similar to $\mathcal{O}$ in B3.2 and using this invariant we prove in 2.22 - 2.25 the completeness in a way similar to the proof of completeness of $[\ ]^1 : \mathrm{PRED} \to \mathcal{I}\mathbf{\Xi}$ in B3.2 - B3.11. Instead of $\vdash_{\mathcal{I}\mathrm{G}}$ we shall mostly write $\vdash$.

Now we repeat Definition B3.2 of the invariant $\mathcal{O}$ for derivations in $\mathcal{I}\mathbf{\Xi}$ for the manysorted structure of our example, in order to contrast it with the one for derivations in $\mathcal{I}\mathbf{G}$ of Definition 2.2 and also because we shall refer to it often in the sequel. Instead of $x_i, \mathcal{V}_i$ and $\mathcal{T}_i$ we shall write $x^{A_i}, \mathcal{V}^{A_i}$ and $\mathcal{T}^{A_i}$.

2.1. Definition (Invariant for derivations in $\mathcal{I}\mathbf{\Xi}$).

$$\begin{aligned}
&\mathcal{T} \equiv \mathcal{T}^{A_1} \mid \mathcal{T}^{A_2};\\
&\mathcal{T}^{A_1} \equiv \mathcal{V}^{A_1} \mid a \mid f\mathcal{T}^{A_1} \mid g\mathcal{T}^{A_1}\mathcal{T}^{A_2};\\
&\mathcal{T}^{A_2} \equiv \mathcal{V}^{A_2};\\
\\
&\mathcal{P} \equiv P\mathcal{T}^{A_1} \mid \mathbf{\Xi}A_i(\lambda x^{A_i}.\mathcal{P}) \mid \mathbf{\Xi}(\mathbf{K}\mathcal{P})(\mathbf{K}\mathcal{P});\\
&\mathcal{G} \equiv \mathbf{L}A_i \mid A_i\mathcal{T}^{A_i} \mid \mathbf{\Xi}A_i(\lambda x^{A_i}.\mathcal{G}) \mid \mathbf{\Xi}(\mathbf{K}\mathcal{P})(\mathbf{K}\mathcal{G}) \mid \mathbf{L}(\mathbf{K}\mathcal{P});\\
&\mathcal{O} \equiv \mathcal{G} \mid \mathcal{P}.
\end{aligned}$$

Now we will define the invariant $\mathcal{O}_G$ for derivations in $\mathcal{I}\mathbf{G}$. $\mathcal{O}_G$ is more complicated then the invariant $\mathcal{O}$ for $\mathcal{I}\mathbf{\Xi}$ for several reasons. If $\Gamma_s^{1,+}, [\Delta]^1, \Gamma(\Delta, \varphi) \vdash \mathbf{L}X$ in system $\mathcal{I}\mathbf{\Xi}$ then $X = A_i$ or $X = \mathbf{K}p$ with $p \in \mathcal{P}$. But if $\Gamma_s^{2,+}, [\Delta]^2, \Gamma(\Delta, \varphi) \vdash \mathbf{L}X$ in system $\mathcal{I}\mathbf{G}$ then we can have for example $X = \mathbf{F}A_1A_2$ or even $X = \mathbf{F}pA_2$. Therefore we define an extension $\mathcal{L}_G$ of the set of sorts $\{A_1, A_2\}$ containing these X's. We need for each $l \in \mathcal{L}_G$ a set of variables $\mathcal{V}_G^l$, similar to the sets $\mathcal{V}^{A_1}$ and $\mathcal{V}^{A_2}$. If we took a different set of variables for each $l \in \mathcal{L}_G$, then the substitution lemma 2.6, that is needed in the proof of Proposition 2.10, would not hold. Therefore we define for each $l \in \mathcal{L}_G$ the skeleton $sk(l)$, such that $sk(p)$ is the same for all $p \in \mathcal{P}_G$ and we define $\mathcal{V}_G^{l_1} = \mathcal{V}_G^{l_2}$ iff $sk(l_1) = sk(l_2)$.

The sets $\mathcal{T}_G^l$ will contain $\lambda-$abstractions: if $l \equiv \mathbf{F}l_1l_2$ and $t^{l_2} \in \mathcal{T}_G^{l_2}$ then $\lambda x^{l_1}.t^{l_2} \in \mathcal{T}_G^l$. This may seem a little bit strange, but in fact it is very natural. For example we have $\vdash \mathbf{F}A_1A_1(\lambda x^{A_1}.x^{A_1})$ by rule $\mathsf{F_i}$. Hence $\lambda x^{A_1}.x^{A_1}$ should be in $\mathcal{T}_G^{\mathbf{F}A_1A_1}$. Moreover in the proof of Lemma 2.9 we really need that $\lambda x^{l_1}.t^{l_2} \in \mathcal{T}_G^{\mathbf{F}l_1l_2}$.

Finally we had $\mathbf{\Xi}(\mathbf{K}p)(\mathbf{K}O) \in \mathcal{O}$ whereas now we have $\mathbf{\Xi}p(\lambda x^p.O) \in \mathcal{O}_G$, where x^p may occur free in O.

2.2. Definition (Invariant $\mathcal{O}_G$ for derivations in $\mathcal{I}\mathbf{G}$). (i) $\Lambda^s(\mathbf{\Xi}, \mathbf{L})$ is, as in Definition B2.12, $\Lambda(\mathbf{\Xi}, \mathbf{L})$ extended by the extra constants A_1, A_2, P, f, g, a associated with the signature s of the manysorted structure of our example.

(ii) Now we will define simultaneously a set $\mathcal{L}_G$ of sorts (an extension of the set $\{A_1, A_2\}$), for each $l \in \mathcal{L}_G$ the skeleton sk(l), a set of variables $\mathcal{V}_G^l$ and a set of terms $\mathcal{T}_G^l$, sets $\mathcal{P}_G, \mathcal{H}_G, \mathcal{G}_G$ and $\mathcal{O}_G$.

Definition of $\mathcal{L}_G$.

$$\mathcal{L}_G \equiv A_1 \mid A_2 \mid Pt^{A_1} \mid \mathbf{G}l_1(\lambda x^{l_1}.l_2) \quad (l_1, l_2 \in \mathcal{L}_G,\ x^{l_1} \in \mathcal{V}_G^{l_1}).$$

Definition of $sk(l)$.

$$\begin{aligned}
&sk(A_i) \equiv A_i;\\
&sk(Pt^{A_1}) \equiv Px^{A_1} \quad \text{(i.e. } sk(Pt^{A_1}) \text{ is independent of } t^{A_1});\\
&sk(\mathbf{G}l_1(\lambda x^{l_1}.l_2)) \equiv \mathbf{G}(sk(l_1))(\lambda x^{l_1}.sk(l_2)).
\end{aligned}$$

Definition of $\mathcal{V}^l_G$.

We assume that for each skeleton $sk(l)$ there is a set of variables

$$\mathcal{V}^{sk(l)}_G \equiv \{x^{sk(l)}, x^{sk(l)}_1, x^{sk(l)}_2, \ldots y^{sk(l)}, y^{sk(l)}_1, y^{sk(l)}_2, \ldots\}$$

such that $\mathcal{V}^{sk(l_1)}_G \cap \mathcal{V}^{sk(l_2)}_G = \emptyset$ if $sk(l_1) \not\equiv sk(l_2)$, and for each $l \in \mathcal{L}_G$ we define

$$\mathcal{V}^l_G \equiv \mathcal{V}^{sk(l)}_G, \quad x^l \equiv x^{sk(l)}, x^l_i \equiv x^{sk(l)}_i, \quad y^l \equiv y^{sk(l)}, y^l_i \equiv y^{sk(l)}_i.$$

Definition of $\mathcal{T}^l_G$.

$$\begin{aligned}
\mathcal{T}^l_G &\equiv a \mid \mathcal{V}^l_G \mid \mathcal{T}^{\mathbf{G}l_1(\lambda x^{l_1}.l)}_G \mathcal{T}^{l_1}_G, && \text{if } l \equiv A_1;\\
\mathcal{T}^l_G &\equiv \quad \mathcal{V}^l_G \mid \mathcal{T}^{\mathbf{G}l_1(\lambda x^{l_1}.l)}_G \mathcal{T}^{l_1}_G, && \text{if } l \equiv A_2 \text{ or } l \equiv Pt^{A_1};\\
\mathcal{T}^l_G &\equiv f \mid \mathcal{V}^l_G \mid \mathcal{T}^{\mathbf{G}l_1(\lambda x^{l_1}.l)}_G \mathcal{T}^{l_1}_G \mid \lambda x^{A_1}.\mathcal{T}^{A_1}_G, && \text{if } l \equiv \mathbf{F}A_1A_1;\\
\mathcal{T}^l_G &\equiv g \mid \mathcal{V}^l_G \mid \mathcal{T}^{\mathbf{G}l_1(\lambda x^{l_1}.l)}_G \mathcal{T}^{l_1}_G \mid \lambda x^{A_1}.\mathcal{T}^{\mathbf{F}A_2A_1}_G, && \text{if } l \equiv \mathbf{F}A_1(\mathbf{F}A_2A_1);\\
\mathcal{T}^l_G &\equiv \quad \mathcal{V}^l_G \mid \mathcal{T}^{\mathbf{G}l_1(\lambda x^{l_1}.l)}_G \mathcal{T}^{l_1}_G \mid \lambda x^{l_2}.\mathcal{T}^{l_3}_G, && \text{if } l \equiv \mathbf{G}l_2(\lambda x^{l_2}.l_3), l \not\in \{\mathbf{F}A_1A_1, \mathbf{F}A_1(\mathbf{F}A_2A_1)\}.
\end{aligned}$$

Definition of $\mathcal{P}_G, \mathcal{H}_G, \mathcal{G}_G$ *and* $\mathcal{O}_G$.

$$\begin{aligned}
&\mathcal{P}_G \equiv P\mathcal{T}^{A_1}_G \mid \mathbf{G}l(\lambda x^l.\mathcal{P}_G);\\
&\mathcal{H}_G \equiv (Pt^{A_1})t^{Pt^{A_1}} \mid \mathbf{\Xi}l(\lambda x^l.\mathcal{H}_G);\\
&\mathcal{G}_G \equiv \mathbf{L}\mathcal{L}_G \mid A_i\mathcal{T}^{A_i}_G \mid \mathbf{\Xi}l(\lambda x^l.\mathcal{G}_G);\\
&\mathcal{O}_G \equiv \mathcal{H}_G \mid \mathcal{G}_G.
\end{aligned}$$

(iii) *Notation.*

$$\begin{aligned}
&l, l_1, l_2, l', \ldots \text{ denote elements of } \mathcal{L}_G;\\
&p, p_1, p_2, p', \ldots \text{ denote elements of } \mathcal{P}_G.
\end{aligned}$$

2.3. Remarks. (i) If $\mathrm{FV}(l_2) = \emptyset$ then $\mathbf{G}l_1(\lambda x^{l_1}.l_2) =_\beta \mathbf{F}l_1l_2$. In that case we will denote $\mathbf{G}l_1(\lambda x^{l_1}.l_2)$ usually by $\mathbf{F}l_1l_2$. In fact we did that already in the above definition.

(ii) The grammar in the definition of $\mathcal{O}_G$ is not context-free and this causes some inaccuracies in the notation. In the definition of $\mathcal{P}_G$ (and similarly at some other places) instead of $\mathbf{G}l(\lambda x^l.\mathcal{P}_G)$ we should write $\mathbf{G}\mathcal{L}_G(\lambda x^{\mathcal{L}_G}.\mathcal{P}_G)$, but then we do not know that these two $\mathcal{L}_G$'s represent the *same* $l \in \mathcal{L}_G$. (We may not have $\mathbf{G}l_1(\lambda x^{l_2}.\mathcal{P}_G)$.)

(iii) In the definition of $\mathcal{T}^l_G$ instead of the clause $\mathcal{T}^{\mathbf{G}l_1(\lambda x^{l_1}.l)}_G \mathcal{T}^{l_1}_G$ one might expect, in view of rule $\mathsf{G_e}$ a clause $t_1^{\mathbf{G}l_1(\lambda x^{l_1}.l')} t_2^{l_1}$ where $l = l'[x^{l_1} := t_2^{l_1}]$, but that does not make a difference because we have for $l \in \mathcal{L}_G$

$$\mathcal{T}^{l[x^{l_1}:=t^{l_1}]}_G = \mathcal{T}^l_G,$$

as follows easily from

$$\mathcal{V}^{l[x^{l_1}:=t^{l_1}]}_G = \mathcal{V}^l_G$$

in Lemma 2.6.

2.4. Lemma. (i) $\mathbf{\Xi} l(\lambda x^l.pt) = \mathbf{G} l(\lambda x^l.p)(\lambda x^l.t)$;
(ii) $\mathbf{G} l(\lambda x^l.p)t = \mathbf{\Xi} l(\lambda x^l.p(tx^l)), \quad x^l \notin \mathrm{FV}(t)$;
(iii) $\mathbf{\Xi} l(\lambda x^l.pt^p) = \mathbf{G} l(\lambda x^l.p) t^{\mathbf{G}l(\lambda x^l.p)}, \quad$ *for* $t^{\mathbf{G}l(\lambda x^l.p)} = \lambda x^l.t^p$;
(iv) $\mathbf{G} l(\lambda x^l.p) t^{\mathbf{G}l(\lambda x^l.p)} = \mathbf{\Xi} l(\lambda x^l.pt^p), \quad$ *for* $t^p = t^{\mathbf{G}l(\lambda x^l.p)} x^l$;
(v) $\mathcal{H}_G = \mathcal{P}_G \mathcal{T}_G^{\mathcal{P}_G}$ *modulo* $\beta-$*equality.*

Proof. (i),(ii) immediately from the definition of $\mathbf{G}$,
(iii), (iv) from (i), (ii),
(v) from (iii) and (iv). □

2.5. Remark. By (v) in the previous lemma we have $\mathcal{H}_G = \mathcal{P}_G \mathcal{T}_G^{\mathcal{P}_G}$. In B3.2 we defined $\mathcal{O} = \mathcal{P} \mid \mathcal{G}$ whereas now we have $\mathcal{O}_G = \mathcal{H}_G \mid \mathcal{G}_G$. This corresponds with the fact that now we treat derivations $\Gamma \vdash [\varphi]^2 M$ instead of $\Gamma \vdash [\varphi]^1$.

2.6. Lemma. *Let* $\mathcal{W} = \mathcal{L}_G, \mathcal{T}^l_G, \mathcal{P}_G, \mathcal{H}_G, \mathcal{G}_G$, *or* $\mathcal{O}_G$ *and let* $w \in \mathcal{W}, x^{l_1} \in \mathcal{V}^{l_1}_G, t^{l_1} \in \mathcal{T}^{l_1}_G$, *then*

$$w[x^{l_1} := t^{l_1}] \in \mathcal{W},$$

moreover if $\mathcal{W} = \mathcal{L}_G$ *then*

$$sk(w[x^{l_1} := t^{l_1}]) \equiv sk(w).$$

and hence

$$\mathcal{V}^{w[x^{l_1}:=t^{l_1}]}_G \equiv \mathcal{V}^w_G.$$

Proof. By induction on the structure of w. Instead of x^{l_1} and t^{l_1} we shall write x and t.

Case $w \in \mathcal{L}_G$.

$w \equiv A_i$. This is trivally ok.
$w \equiv P t_1^{A_1}$. This follows from the case $w \equiv t_1^{A_1}$.
$w \equiv \mathbf{G} l_2(\lambda y^{l_2}.l_3)$. Then $w[x := t] \equiv \mathbf{G}(l_2[x := t])(\lambda y^{l_2}.l_3[x := t]) \equiv$
$\equiv \mathbf{G}(l_2[x := t])(\lambda y^{l_2[x:=t]}.l_3[x := t])$
(by the second part of the IH for l_2) and this is in $\mathcal{L}_G$ by the first part of the IH for l_2 and l_3. The second part of the IH for l_2 and l_3 yields $sk(w[x := t]) \equiv sk(w)$.

Case $w = t^l \in \mathcal{T}_G^l$.

$l \equiv A_1$. Now $w \equiv x_1^{A_1}, a$ or $t_1^{\mathbf{G}l_1(\lambda x_1^{l_1}.A_1)} t_2^l$. All these cases are easy.
$l \equiv A_2$ or $Pt_1^{A_1}$. Easy.
$l \equiv \mathbf{G}l_2(\lambda y^{l_2}.l_3)$. Now $w \equiv x_1^l, t_1^{\mathbf{G}l_4(\lambda x_1^{l_4}.l)} t_2^{l_4}, \lambda y^{l_2}.t_1^{l_3}, f$ or g. Easy again.

Case $w \in \mathcal{P}_G$.

$w \equiv Pt_1^{A_1}$. This follows from the case $w \equiv t_1^{A_1}$.
$w \equiv \mathbf{G}l_2(\lambda y^{l_2}.p)$. Then $w[x := t] \equiv \mathbf{G}(l_2[x := t])(\lambda y^{l_2}.p[x := t]) \equiv$
$\equiv \mathbf{G}(l_2[x := t])(\lambda y^{l_2[x:=t]}.p[x := t])$.

Cases $w \in \mathcal{H}_G, w \in \mathcal{G}_G$. Similarly. □

2.7. Lemma. *Let* $cX_1 \dots X_n =_{\beta\eta} M$ *for some constant* c *and some* $M \in \mathcal{O}_G$. *Then* $n \in \{1, 2\}$ *and* $M \equiv cY_1 \dots Y_n$ *with* $Y_i =_{\beta\eta} X_i$.

Proof. By Church-Rosser for $\beta\eta$-reduction. □

In the definition of $\mathcal{G}_G$ one of the clauses is $A_i\mathcal{T}_G^{A_i}$ where one might expect the more general clause $l\mathcal{T}_G^l$. But $A_i\mathcal{T}_G^{A_i}$ is sufficient as follows from the next lemma.

2.8. Lemma.

$$lt^l \in \overline{\mathcal{O}_G} \text{ for each } l \in \mathcal{L}_G.$$

Proof. By induction on the structure of l. It is clear if $l = A_1,\ A_2$ or Pt^{A_1}. So let $l = \mathbf{G}l_1(\lambda x^{l_1}.l_2)$. Then $lt^l = \mathbf{G}l_1(\lambda x^{l_1}.l_2)t^l = \mathbf{\Xi}l_1(\mathbf{S}(\lambda x^{l_1}.l_2)t^l) = \mathbf{\Xi}l_1(\lambda x^{l_1}.l_2(t^l x^{l_1}))$. Now $t^l \in \mathcal{T}_G^l = \mathcal{T}_G^{\mathbf{G}l_1(\lambda x^{l_1}.l_2)}$ and $x^{l_1} \in \mathcal{T}_G^{l_1}$ hence $t^l x^{l_1} \in \mathcal{T}_G^{l_2}$. So by the induction hypothesis one has $l_2(t^l x^{l_1}) \in \overline{\mathcal{G}_G}$. Hence $\mathbf{\Xi}l_1(\lambda x^{l_1}.l_2(t^l x^{l_1})) \in \overline{\mathcal{G}_G}$. □

2.9. Lemma.

$$lV \in \overline{\mathcal{O}_G} \ \Rightarrow\ V \in \overline{\mathcal{T}_G^l}.$$

Proof. By induction on the structure of l. It is clear if $l = A_1, A_2$ or Pt^{A_1}. So let $l = \mathbf{G}l_1(\lambda x^{l_1}.l_2)$. Then $lV = \mathbf{\Xi}l_1(\lambda x^{l_1}.l_2(Vx^{l_1}))$. So $l_2(Vx^{l_1}) \in \overline{\mathcal{O}_G}$ and hence by the induction hypothesis we have $Vx^{l_1} = t^{l_2} \in \mathcal{T}_G^{l_2}$. So $V = \lambda x^{l_1}.t^{l_2} \in \mathcal{T}_G^l$. □

Now we will prove $\Gamma \vdash M, \Gamma \subseteq \overline{\mathcal{O}_G} \ \Rightarrow\ M \in \overline{\mathcal{O}_G}$. The proof is similar to the proof of Proposition B3.10, but without the induction loading. We used that induction loading to derive, in subcase Ξ_{i}(a), that if $\Gamma \vdash \mathbf{\Xi}(\mathbf{K}p)(\lambda x.O)$ then $x \notin \mathrm{FV}(O)$. But now we have $\Gamma \vdash \mathbf{\Xi}(p)(\lambda x.O)$ where x may occur free in O.

2.10. Proposition. *$\mathcal{O}_G$ is an invariant (modulo $\beta\eta-$equality) for derivations in $\mathcal{I}\mathbf{G}$:*

$$\Gamma \vdash M, \Gamma \subseteq \overline{\mathcal{O}_G} \ \Rightarrow\ M \in \overline{\mathcal{O}_G}.$$

PROOF. By induction on the derivation of $\Gamma \vdash M$. We only consider the three specific rules $\Xi_{\mathrm{e}}, \Xi_{\mathrm{i}}$ and G_{L}. (The two general rules are easy and $\mathsf{G}_{\mathrm{e}}, \mathsf{G}_{\mathrm{i}}$ are equivalent to $\Xi_{\mathrm{e}}, \Xi_{\mathrm{i}}$.)

Case Ξ_{e}. $\Gamma \vdash M$ is $\Gamma \vdash YV$ as direct consequence of $\Gamma \vdash \mathbf{\Xi}XY$, $\Gamma \vdash XV$.

By the IH one has $\mathbf{\Xi}XY \in \overline{\mathcal{O}_G}, XV \in \overline{\mathcal{O}_G}$. From $\mathbf{\Xi}XY \in \overline{\mathcal{O}_G}$ it follows that $X = l, Y = \lambda x^l.O$ where $O \in \mathcal{O}_G$. Now $XV = lV \in \overline{\mathcal{O}_G}$, hence $V = t^l \in \mathcal{T}_G^l$ by Lemma 2.9. So $M = (\lambda x^l.O)t^l = O[x^l := t^l] \in \overline{\mathcal{O}_G}$ by Lemma 2.6.

Case Ξ_{i}. $\Gamma \vdash M$ is $\Gamma \vdash \mathbf{\Xi}XY$ as direct consequence of
$\Gamma \vdash \mathbf{L}X$, $\Gamma, Xx \vdash Yx$ with $x \notin \mathrm{FV}(\Gamma, X, Y)$.

By the IH one has $\mathbf{L}X \in \overline{\mathcal{O}_G}$, hence $X = l \in \mathcal{L}_G$. Now x is any variable, so we may assume that $x = x^l \in \mathcal{V}_G^l$. Then $lx \in \overline{\mathcal{O}_G}$ by Lemma 2.8, hence by the IH we have $Yx \in \overline{\mathcal{O}_G}$. Let $Yx = O \in \overline{\mathcal{O}_G}$. Then $M = \mathbf{\Xi}l(\lambda x^l.O) \in \overline{\mathcal{O}_G}$.

Case G_{L}. $\Gamma \vdash M$ is $\Gamma \vdash \mathbf{L}(\mathbf{G}XY)$ as direct consequence of
$\Gamma \vdash \mathbf{L}X$, $\Gamma, Xx \vdash \mathbf{L}(Yx)$ with $x \notin \mathrm{FV}(\Gamma, X, Y)$.

By the IH one has $\mathbf{L}X \in \overline{\mathcal{O}_G}$, hence $X = l \in \mathcal{L}_G$. Again we may assume that $x = x^l \in \mathcal{V}_G^l$. Then $lx \in \overline{\mathcal{O}_G}$, hence by the IH we have $\mathbf{L}(Yx) \in \overline{\mathcal{O}_G}$. So $Yx = l_1 \in \mathcal{L}_G$. We get $\mathbf{G}XY = \mathbf{G}l(\lambda x^l.l_1)) \in \overline{\mathcal{L}_G}$. Hence $\mathbf{L}(\mathbf{G}XY) \in \overline{\mathcal{G}_G} \subseteq \overline{\mathcal{O}_G}$. □

The invariant $\mathcal{O}_G$ is more complicated than $\mathcal{O}$ and we cannot prove completeness directly using this invariant, as in in B3.11 for $\mathcal{I}\mathbf{\Xi}$ relative to PRED. We will show in 2.11-2.20 that in each proof $\Gamma \vdash_{\mathcal{I}\mathsf{G}} M$, where $\Gamma \subseteq \overline{\mathcal{O}_G}$, all cut formulas can be eliminated. Then using another invariant $\mathcal{O}_1$, similar to $\mathcal{O}$ we prove in Proposition 2.25 completeness of $\mathcal{I}\mathbf{G}$ for PRED in a way similar to the proof of completeness of $\mathcal{I}\mathbf{\Xi}$ for PRED.

Cut elimination

First we define the $\mathbf{\Xi}$–length $L_{\mathbf{\Xi}}(O)$ for $O \in \overline{\mathcal{O}_G}$ (and $L_{\mathbf{\Xi}}(l)$ for $l \in \mathcal{L}_G$) and state some lemma's, without the (trivial) proofs.

2.11. DEFINITION ($\mathbf{\Xi}$–length $L_{\mathbf{\Xi}}$).

$$L_{\mathbf{\Xi}}((Pt^{A_1})t^{Pt^{A_1}}) = L_{\mathbf{\Xi}}(\mathbf{L}l) = L_{\mathbf{\Xi}}(A_i t^{A_i}) = 0, \quad L_{\mathbf{\Xi}}(\mathbf{\Xi}l(\lambda x^l.O)) = 1 + L_{\mathbf{\Xi}}(l) + L_{\mathbf{\Xi}}(O),$$

$$L_{\mathbf{\Xi}}(A_i) = L_{\mathbf{\Xi}}(Pt^{A_1}) = 0, \quad L_{\mathbf{\Xi}}(\mathbf{G}l(\lambda x^l.q)) = 1 + L_{\mathbf{\Xi}}(l) + L_{\mathbf{\Xi}}(q).$$

2.12. LEMMA. (i) $X, Y \in \mathcal{L}_G \cup \mathcal{O}_G$, $X \twoheadrightarrow_{\beta\eta} Y \Rightarrow L_{\mathbf{\Xi}}(X) = L_{\mathbf{\Xi}}(Y)$.
(ii) $X, Y \in \mathcal{L}_G \cup \mathcal{O}_G$, $X =_{\beta\eta} Y \Rightarrow L_{\mathbf{\Xi}}(X) = L_{\mathbf{\Xi}}(Y)$.

2.13. DEFINITION. Let $M \in \overline{\mathcal{O}_G}$, $M =_{\beta\eta} O$. Then $L_{\mathbf{\Xi}}(M) = L_{\mathbf{\Xi}}(O)$.

2.14. Lemma. $L_{\Xi}(lt^l) = L_{\Xi}(l)$.

2.15. Lemma. $X \in \mathcal{L}_G \cup \overline{\mathcal{O}_G},\ t^l \in \mathcal{T}_G^l \ \Rightarrow\ L_{\Xi}(X[x^l := t^l]) = L_{\Xi}(X)$.

2.16. Lemma.

$$\Xi XY \in \overline{\mathcal{O}_G},\ XV \in \overline{\mathcal{O}_G} \ \Rightarrow\ L_{\Xi}(XV) < L_{\Xi}(\Xi XY)\ \&\ L_{\Xi}(YV) < L_{\Xi}(\Xi XY).$$

Proof. $\Xi XY = \Xi l(\lambda x^l.O)$ and $V \in \overline{\mathcal{T}_G^l}$. Now use Lemma's 2.14 and 2.15. □

2.17. Definition. A cut elimination is a proof reduction of the following form



ΞXY is called the cut formula.

2.18. Lemma. *In a proof of* $\Gamma \vdash_{\mathcal{I}G} M$ *(or* $\Gamma \vdash_{\mathcal{I}\Xi} M$ *), where* $\Gamma \subseteq \overline{\mathcal{O}_G}$, *all cut formulas can be eliminated by the above proof reduction.*

Proof. By induction on the maximal Ξ−length of cut formulas in the proof of $\Gamma \vdash_{\mathcal{I}G} M$. Consider a cut in the proof with a cut formula ΞXY of maximal Ξ−length. Eliminate this cut. Now new cut formulas may have been introduced at XV and YV, however by Lemma 2.16 these have shorter Ξ−length. In the same way all other cut formulas of maximal Ξ−length can be eliminated and so by the induction hypothesis all cut formulas can be eliminated. □

2.19. Remark. The above can easily be generalised to allow $\beta\eta$−equality steps (Eq-steps) as follows.



2.20. REMARK. Similarly we can define cut elimination for $\mathcal{IP}$ as a proof reduction of the following form.



This is in fact a special case of Definition 2.17. (Substitute $\mathbf{K}X$ for X and $\mathbf{K}Y$ for Y in Definition 2.17.) Also in a proof of $\Gamma \vdash_{\mathcal{IP}} M$ where $\Gamma \subseteq \overline{\mathcal{O}_G}$ all cut formulas can be eliminated.

Now we define similar to the set $\mathcal{O}$ in B3.2 a set $\mathcal{O}_1$ such that $N \subseteq \overline{\mathcal{O}_1}$ for terms N of interest occurring in a cut free proof of $\Gamma_s^{2,+}, [\Delta]^2, \Gamma(\Delta, \varphi) \vdash_{\mathcal{IG}} [\varphi]^2 M$.

2.21. DEFINITION. *Definition of* $\mathcal{L}_1$.

$$\mathcal{L}_1 \equiv A_1 \mid A_2 \mid \mathcal{P}_1.$$

Definition of $sk(l)$ *and* $\mathcal{V}_1^l$.

$$\begin{aligned}
&sk(A_i) \equiv A_i;\\
&sk(Pt^{A_1}) \equiv Px^{A_1} \quad \text{(i.e. } sk(Pt^{A_1}) \text{ is independent of } t^{A_1}\text{)};\\
&sk(\mathbf{G}l_2(\lambda x^{l_2}.p_1)) \equiv \mathbf{G}(sk(l_2))(\lambda x^{l_2}.sk(p_1)).
\end{aligned}$$

$$\mathcal{V}_1^l \equiv \{x_1^p, x_2^p, \ldots\} \quad , x_i^l = x_i^{sk(l)}, x_i^{p_1} \neq x_i^{p_2} \text{ if } sk(p_1) \not\equiv sk(p_2);$$

Definition of $\mathcal{T}_1^l$.

$$\begin{aligned}
&\mathcal{T}_1^l \equiv \mathcal{V}_1^{A_1} \mid a \mid f\mathcal{T}_1^{A_1} \mid g\mathcal{T}_1^{A_1}\mathcal{T}_1^{A_2} && \text{if } l \equiv A_1;\\
&\mathcal{T}_1^l \equiv \mathcal{V}_1^{A_2} && \text{if } l \equiv A_2;\\
&\mathcal{T}_1^l \equiv \mathcal{V}_1^l \mid \mathcal{T}_1^{\mathbf{G}l_1(\lambda x^{l_1}.l)}\mathcal{T}_1^{l_1} && \text{if } l \equiv Pt^{A_1};\\
&\mathcal{T}_1^l \equiv \mathcal{V}_1^l \mid \mathcal{T}_1^{\mathbf{G}l_1(\lambda x^{l_1}.l)}\mathcal{T}_1^{l_1} \mid \lambda x^{l_2}.\mathcal{T}_1^{p_1}, && \text{if } l \equiv \mathbf{G}l_2(\lambda x^{l_2}.p_1)).
\end{aligned}$$

Definition of $\mathcal{P}_1, \mathcal{H}_1, \mathcal{G}_1$ *and* $\mathcal{O}_1$.

$$\begin{aligned}
&\mathcal{P}_1 \equiv P\mathcal{T}_1^{A_1} \mid \mathbf{G}l(\lambda x^l.\mathcal{P}_1);\\
&\mathcal{H}_1 \equiv (Pt^{A_1})t^{Pt^{A_1}} \mid \mathbf{\Xi}l(\lambda x^l.\mathcal{H}_1);\\
&\mathcal{G}_1 \equiv \mathbf{L}\mathcal{L}_1 \mid A_i\mathcal{T}_1^{A_i} \mid \mathbf{\Xi}l(\lambda x^l.\mathcal{G}_1);\\
&\mathcal{O}_1 \equiv \mathcal{H}_1 \mid \mathcal{G}_1.
\end{aligned}$$

2.22. Remarks. (i) Similar to Lemma 2.4(v) we now have $\mathcal{H}_1 = \mathcal{P}_1\mathcal{T}_1^{\mathcal{P}_1}$.

(ii) If $l \equiv A_i$ in $\mathbf{G}l(\lambda x^l.p_1)$ in the definition of $\mathcal{P}_1$, then x^l may occur free in p_1; but if $l \equiv p$ then $x^l \notin \mathrm{FV}(p_1)$. So $\mathbf{G}l(\lambda x^l.p_1)$ may be split into $\mathbf{G}A_i(\lambda x^{A_i}.p_1)$ and $\mathbf{G}p(\mathbf{K}p_1)$ and similarly for $\mathbf{\Xi}l(\lambda x^l.g)$ in the definition of $\mathcal{G}_1$.

Lemma's 2.6 - 2.9 mutatis mutandis also hold for $\mathcal{O}_1$. Moreover lemma's corresponding to Lemma's B3.7 - B3.9 hold. Replace $[-]^1$ by $[-]^2$ and in the lemma corresponding to B3.7 take $\mathcal{H}_1$ instead of $\mathcal{P}_1$.

2.23. Lemma. *Let $p, p' \in \mathcal{P}_1$. Then $pM = p'M' \;\Rightarrow\; p = p'$.*

Proof. Induction on p. □

The following lemma corresponds to Proposition B3.10.

2.24. Lemma. *Let $\Gamma \subseteq \overline{\mathcal{O}_1}$ and let $\Gamma \vdash_{\mathcal{I}\mathrm{G}} N$ be a cut free proof. Then*

(i) *If the last applied rule in the proof is not the $\Xi_{\mathrm{i}}-$rule, nor the $\mathsf{G}_{\mathsf{L}}-$rule (modulo the $\beta\eta-$rule), then $N \in \overline{\mathcal{O}_1}$.*

(ii) *If $N = lt$ with $l \in \mathcal{L}_1, t \in \mathcal{T}_G^l$ then $N \in \overline{\mathcal{O}_1}$.*

Proof. The proof of (i) and (ii) is by simultaneous induction on $\Gamma \vdash N$. The two general cases are easy, so we only consider Cases Ξ_{i}, for (ii), and Ξ_{e}.

Case Ξ_{i}. $\Gamma \vdash N$ is $\Gamma \vdash lt = \mathbf{\Xi}XY$ as direct consequence of $\Gamma \vdash \mathbf{L}X$, $\Gamma, Xx \vdash Yx$.

Now $lt = \mathbf{G}l_1(\lambda x^{l_1}.p)t = \mathbf{\Xi}l_1(\lambda x^{l_1}.p(tx^{l_1})) = \mathbf{\Xi}XY$. The IH for $\Gamma, l_1x^{l_1} \vdash p(tx^{l_1})$ yields $p(tx^{l_1}) = O \in \overline{\mathcal{O}_1}$. So $\mathbf{\Xi}l_1(\lambda x^{l_1}.p(tx^{l_1})) = \mathbf{\Xi}l_1(\lambda x^{l_1}.O) \in \overline{\mathcal{O}_1}$.

Case Ξ_{e}. $\Gamma \vdash N$ is $\Gamma \vdash YV$ as direct consequence of $\Gamma \vdash \mathbf{\Xi}XY$, $\Gamma \vdash XV$.

By the induction hypothesis for $\Gamma \vdash \mathbf{\Xi}XY$ we have $\mathbf{\Xi}XY \in \overline{\mathcal{O}_1}$. So $\mathbf{\Xi}XY = \mathbf{\Xi}l_1(\lambda x^{l_1}.O)$ The IH for $\Gamma \vdash l_1V$ yields $l_1V \in \overline{\mathcal{O}_1}$, hence $V \in \overline{\mathcal{T}_1^{l_1}}$. So $YV = O[x^{l_1} := V] \in \overline{\mathcal{O}_1}$. □

2.25. Proposition (Completeness of $\mathcal{I}\mathbf{G}$ for PRED).

$$\Gamma_s^{2,+}, [\Delta]^2, \Gamma(\Delta, \varphi) \vdash_{\mathcal{I}\mathrm{G}} [\varphi]^2 M \;\Rightarrow\; \Delta \vdash_{\mathrm{PRED}} \varphi.$$

Proof. The proof resembles very much the proof of Proposition B3.11.
Note that $\Gamma_s^{2,+} \subseteq \overline{\mathcal{G}_1}$ and $\Gamma(\Delta, \varphi) \subseteq \overline{\mathcal{G}_1}$. Hence it is sufficient to prove

$$\Gamma_g, [\Delta]^2 \vdash_{\mathcal{I}\mathrm{G}} N,\;\; \Gamma_g \subseteq \overline{\mathcal{G}_1},\;\; N = [\varphi]^2 M \;\;\Rightarrow\;\; \Delta \vdash_{\mathrm{PRED}} \varphi. \qquad (**)$$

Write $\Gamma = \Gamma_g, [\Delta]^2$. Then $\Gamma \subseteq \overline{\mathcal{O}_1}$. The proof of $(**)$ is by induction on the derivation of $\Gamma \vdash_{\mathcal{I}\mathrm{G}} N$. We may assume that the derivation is cut free.

Case 1. $\Gamma \vdash N$ because $N \in \Gamma$.

$N = [\varphi]^2 M$, so $N \notin \overline{\mathcal{G}_1}$, hence $N \in [\Delta]^2$. We get from Lemma 2.23 that $[\varphi]^2 = [\psi]^2$ for some $\psi \in \Delta$. so $\varphi = \psi$ and hence $\varphi \in \Delta$. Therefore $\Delta \vdash_{\mathrm{PRED}} \varphi$.

Case 2. $\Gamma \vdash N$ is a direct consequence of $\Gamma \vdash N'$ and $N = N'$.

Now $N' = N = [\varphi]^2 M$ and by the IH for N' one has $\Delta \vdash_{\mathrm{PRED}} \varphi$.

Case Ξ_{e}. $\Gamma \vdash N$ is $\Gamma \vdash YV$ as direct consequence of $\Gamma \vdash \mathbf{\Xi} XY$, $\Gamma \vdash XV$.

$\Gamma \vdash N$ is cut free, hence from Lemma 2.24 we get that $\mathbf{\Xi} XY \in \overline{\mathcal{O}_1}$. We distinguish 3 subcases.

Subcase Ξ_{e}(a). $X = A_i, Y = \lambda x^{A_i}.h$. Now $h = pt^p = [\psi]^2 t^p$ for some $p \in \mathcal{P}_1, \psi \in \mathbb{F}_{\mathrm{PRED}}$. We have $XV = A_i V \in \overline{\mathcal{O}_1}$ by Lemma 2.24. Therefore $V = t^{A_i} = [t_{\mathbf{A}_i}]^2$. Now $N = [\varphi]^2 M = YV = (\lambda x^{A_i}.[\psi]^2 t^p)[t_{\mathbf{A}_i}]^2 = [\psi]^2[x^{A_i} := [t_{\mathbf{A}_i}]^2]t' = [\psi[x^{\mathbf{A}_i} := t_{\mathbf{A}_i}]]^2 t'$ for some t'. So $[\varphi]^2 \equiv [\psi[x^{\mathbf{A}_i} := t_{\mathbf{A}_i}]]^2$ Hence $\varphi \equiv \psi[x^{\mathbf{A}_i} := t_{\mathbf{A}_i}]$. $\mathbf{\Xi} XY = \mathbf{\Xi} A_i(\lambda x^{A_i}.pt^p) = \mathbf{G} A_i(\lambda x^{A_i}.p)t'$ (for some t') $= [\forall x^{\mathbf{A}_i} \psi]^2 t'$. So we have by the IH $\Delta \vdash_{\mathrm{PRED}} \forall x^{\mathbf{A}_i} \psi$. Hence $\Delta \vdash_{\mathrm{PRED}} \psi[x^{\mathbf{A}_i} := t_{\mathbf{A}_i}] \equiv \varphi$.

Subcase Ξ_{e}(b). $X = p, Y = \lambda x^p.h$. Now $Y = \lambda x^p.p_2 t^{p_2}$ where $x^p \notin \mathrm{FV}(p_2)$. Then $N = [\varphi]^2 M = YV = p_2 t'$ for some t'. So $[\varphi]^2 = p_2$. Let $p = [\varphi_1]^2$. Then $\mathbf{\Xi} XY = \mathbf{G} p(\mathbf{K} p_2) t'' = \mathbf{F} p p_2 t'' = [\varphi_1 \supset \varphi]^2 t''$ for some t''. So we have by the IH

$$\Delta \vdash_{\mathrm{PRED}} \varphi_1 \supset \varphi.$$

Also $\Gamma \vdash XV = [\varphi_1]^2 V$. So by the IH one has

$$\Delta \vdash_{\mathrm{PRED}} \varphi_1.$$

Therefore it follows by modus ponens that

$$\Delta \vdash_{\mathrm{PRED}} \varphi.$$

Subcase Ξ_{e}(c). $\mathbf{\Xi} XY = \mathbf{\Xi} l(\lambda x^l.g.)$ This case is not applicable because $YV \neq [\varphi]^2 M$.

Case Ξ_{i}. $\Gamma \vdash N$ is $\Gamma \vdash \mathbf{\Xi} XY$ as direct consequence of
$\Gamma \vdash \mathbf{L} X$, $\Gamma, Xx \vdash Yx$ with $x \notin \mathrm{FV}(\Gamma, X, Y)$.

$\mathbf{\Xi} XY = [\varphi]^2 M$. So $[\varphi]^2 = \mathbf{G} l_1(\lambda x^{l_1}.p_2)$ for some $l_1 \in \mathcal{L}_1$ and $p_2 \in \mathcal{P}_1$, and $\mathbf{\Xi} XY = \mathbf{\Xi} l_1(\lambda x^{l_1}.p_2(Mx^{l_1}))$. We consider 2 subcases.

Subcase Ξ_{i}(a). $X = A_i$. Let $p_2 = [\psi]^2$. Then $\varphi = \forall x^{\mathbf{A}_i} \psi$. As $\Gamma, Xx^{A_i} \vdash p_2(Mx^{A_i})$ one has $\Delta \vdash_{\mathrm{PRED}} \psi$ by the IH. Now x^{A_i} does not occur in Γ, so

$$\Delta \vdash_{\mathrm{PRED}} \forall x^{\mathbf{A}_i} \psi.$$

Subcase Ξ_{i}(b). $X = p_1$. Let $p_1 = [\varphi_1]^2, p_2 = [\varphi_2]^2$. Then $\varphi = \varphi_1 \supset \varphi_2$. Now $\Gamma, [\varphi_1]^2 x \vdash [\varphi_2]^2(Mx)$. So one has by the IH $\Delta, \varphi_1 \vdash_{\mathrm{PRED}} \varphi_2$. Hence

$$\Delta \vdash_{\mathrm{PRED}} \varphi_1 \supset \varphi_2.$$

Case $\mathsf{G_L}$. $\Gamma \vdash N$ is $\Gamma \vdash \mathbf{L}(\mathbf{G}XYZ)$ as direct consequence of $\Gamma \vdash \mathbf{L}X$, $\Gamma, Xx \vdash \mathbf{L}(Yx)$ with $x \notin \mathrm{FV}(\Gamma, X, Y)$.

This case is not applicable because $\mathbf{L}(\mathbf{G}XYZ) \neq [\varphi]^2 M$. $\square$

3. Completeness of $\mathcal{IF}$ for PROP

We must show for $\varphi \in \mathbb{F}_{\mathrm{PROP}}$, $\Delta \subseteq \mathbb{F}_{\mathrm{PROP}}$:

$$\forall r \exists M \in \Lambda [[\Delta]^2_r, \Gamma^2_r(\Delta, \varphi) \vdash_{\mathcal{IF}} [\varphi]^2_r M] \;\Rightarrow\; \Delta \vdash_{\mathrm{PROP}} \varphi.$$

It is sufficient to show

$$\exists M \in \Lambda [[\Delta]^2_{\mathsf{I}}, \Gamma^2_{\mathsf{I}}(\Delta, \varphi) \vdash_{\mathcal{IF}} [\varphi]^2_{\mathsf{I}} M] \;\Rightarrow\; \Delta \vdash_{\mathrm{PROP}} \varphi.$$

We omit I in the notation and we prove

$$[\Delta]^2, \Gamma^2(\Delta, \varphi) \vdash_{\mathcal{IF}} [\varphi]^2 M \;\Rightarrow\; \Delta \vdash_{\mathrm{PROP}} \varphi.$$

The proof is much simpler than the proof of completeness of $\mathcal{IG}$ for PRED, because there is no need for cut elimination. We define only one invariant $\mathcal{O}_2$ in 3.1, similar to $\mathcal{O}$ and $\mathcal{O}_1$ and using this invariant we prove the completeness in 3.2 - 3.5.

3.1. Definition (Invariant for derivations in $\mathcal{IF}$). $\mathcal{V}_2$ is a set of proposition variables.
For every $p \in \mathcal{P}_2$ there is a different set $\mathcal{V}_2^p$ of variables.

Definition of $\mathcal{T}_2^p$.

$$\begin{aligned} \mathcal{T}_2^p &\equiv \mathcal{V}_2^p \mid \mathcal{T}_2^{\mathbf{F}p_1 p} \mathcal{T}_2^{p_1}, && \text{if } p \in \mathcal{V}_2; \\ \mathcal{T}_2^p &\equiv \mathcal{V}_2^p \mid \mathcal{T}_2^{\mathbf{F}p_1 p} \mathcal{T}_2^{p_1} \mid \lambda x^{p_2}.\mathcal{T}_2^{p_3}, && \text{if } p \equiv \mathbf{F}p_2 p_3. \end{aligned}$$

Definition of $\mathcal{P}_2, \mathcal{H}_2, \mathcal{G}_2$ *and* $\mathcal{O}_2$.

$$\begin{aligned} &\mathcal{P}_2 \equiv \mathcal{V}_2 \mid \mathbf{F}\mathcal{P}_2\mathcal{P}_2; \\ &\mathcal{H}_2 \equiv vt^v (v \in \mathcal{V}_2) \mid \mathbf{\Xi} p_1 (\lambda x^{p_1}.\mathcal{H}_2); \\ &\mathcal{G}_2 \equiv \mathbf{L}\mathcal{P}_2 \mid \mathbf{\Xi}\mathcal{P}_2(\mathbf{K}\mathcal{G}_2); \\ &\mathcal{O}_2 \equiv \mathcal{H}_2 \mid \mathcal{G}_2. \end{aligned}$$

3.2. Remark. Similarly to Lemma 2.4(v) we have $\mathcal{H}_2 = \mathcal{P}_2 \mathcal{T}_2^{\mathcal{P}_2}$. Note that in $\mathbf{\Xi} p_1 (\lambda x^{p_1}.h)$ in the definition of $\mathcal{H}_2$ we have $h = p_2 t^{p_2}$ where $x^{p_1} \notin \mathrm{FV}(p_2)$.

We have lemma's similar to Lemma's 2.6 - 2.9, with similar, but easier, proofs. (Now $\mathcal{L}_2 = \mathcal{P}_2$.)

3.3. PROPOSITION. *In* $\mathcal{I}\mathbf{F}$ *we have*

$$\Gamma \vdash M, \Gamma \subseteq \overline{\mathcal{O}_2} \;\Rightarrow\; M \in \overline{\mathcal{O}_2}.$$

PROOF. Similar to the proof of B3.10. Instead of *Case* G_L we now have *Case* F_L. □

3.4. LEMMA. (i) $\mathcal{G}_2 \cap \mathcal{H}_2 = \emptyset$;
(ii) $\overline{\mathcal{G}_2} \cap \overline{\mathcal{G}_2} = \emptyset$;
(iii) $[-]^2 : \mathbb{F}_{\mathrm{PROP}} \to \mathcal{P}_2$ *is a bijection.*
(iv) *Let* $p, p' \in \mathcal{P}_2$. *Then* $pM = p'M' \;\Rightarrow\; p = p'$.

PROOF. By easy inductions. For (ii) we use Church Rosser. □

3.5. PROPOSITION (Completeness of $\mathcal{I}\mathbf{F}$ for PROP).

$$[\Delta]^2, \Gamma^2(\Delta, \varphi) \vdash_{\mathcal{I}\mathrm{F}} [\varphi]^2 \;\Rightarrow\; \Delta \vdash_{\mathrm{PROP}} \varphi.$$

PROOF. Similar to the proof of Proposition 2.25 but easier. The subcases corresponding to subcases Ξ_{e}(a) and Ξ_{i}(a) do not occur. □

4. Completeness of $\mathcal{I}\mathbf{G}$ for PRED$\lambda\to$

In this section we will show completeness of $\mathcal{I}\mathbf{G}$ over predicate logic extended with higher type functions. In this logic we may quantify over all functions based on $\mathbf{A}_1$ and $\mathbf{A}_2$ and also over functions ranging over functions etc. and we can state e.g.

$$\forall f_1 \in ((\mathbf{A}_1 \to \mathbf{A}_2) \to \mathbf{A}_1) \forall f_2 \in (\mathbf{A}_1 \to \mathbf{A}_2)[\mathbf{P}(f_1 f_2)].$$

4.1. DEFINITION. PRED$\lambda\to_0$, a preliminary version of Predicate Calculus with Higher Type Functions, is defined by

(i) *Types of* PRED$\lambda\to_0$

$$\mathbb{L}_{\mathrm{PRED}\lambda\to_0} = \mathbf{A}_1 \mid \mathbf{A}_2 \mid \mathbb{L}_{\mathrm{PRED}\lambda\to_0} \to \mathbb{L}_{\mathrm{PRED}\lambda\to_0}.$$

(ii) *Variables of* PRED$\lambda\to_0$

$$\mathbb{V}^{\mathbf{l}} = \{x_1^{\mathbf{l}}, x_2^{\mathbf{l}}, \ldots\} \quad \text{for } \mathbf{l} \in \mathbb{L}_{\mathrm{PRED}\lambda\to_0}.$$

(iii) *Terms of* PRED$\lambda\to_0$

$$\begin{aligned}
\mathbb{T}^{\mathbf{l}} &= \mathbf{a} \mid \mathbb{V}^{\mathbf{l}} \mid \mathbb{T}^{\mathbf{l}_1 \to \mathbf{l}} \mathbb{T}^{\mathbf{l}_1}, && \text{if } \mathbf{l} = \mathbf{A}_1;\\
\mathbb{T}^{\mathbf{l}} &= \quad \mathbb{V}^{\mathbf{l}} \mid \mathbb{T}^{\mathbf{l}_1 \to \mathbf{l}} \mathbb{T}^{\mathbf{l}_1}, && \text{if } \mathbf{l} = \mathbf{A}_2;\\
\mathbb{T}^{\mathbf{l}} &= \mathbf{f} \mid \mathbb{V}^{\mathbf{l}} \mid \mathbb{T}^{\mathbf{l}_1 \to \mathbf{l}} \mathbb{T}^{\mathbf{l}_1}, && \text{if } \mathbf{l} = \mathbf{A}_1 \to \mathbf{A}_1;\\
\mathbb{T}^{\mathbf{l}} &= \mathbf{g} \mid \mathbb{V}^{\mathbf{l}} \mid \mathbb{T}^{\mathbf{l}_1 \to \mathbf{l}} \mathbb{T}^{\mathbf{l}_1}, && \text{if } \mathbf{l} = \mathbf{A}_1 \to (\mathbf{A}_2 \to \mathbf{A}_1);\\
\mathbb{T}^{\mathbf{l}} &= \quad \mathbb{V}^{\mathbf{l}} \mid \mathbb{T}^{\mathbf{l}_1 \to \mathbf{l}} \mathbb{T}^{\mathbf{l}_1}, && \text{if } \mathbf{l} \notin \{\mathbf{A}_1, \mathbf{A}_2, \mathbf{A}_1 \to \mathbf{A}_1, \mathbf{A}_1 \to (\mathbf{A}_2 \to \mathbf{A}_1)\}.
\end{aligned}$$

(iv) *Well formed formulas of* PRED$\lambda\!\rightarrow_0$

$$\mathbb{F}_{\mathrm{PRED}\lambda\rightarrow_0} = \mathbf{P}\mathbb{T}^{\mathbf{A}_1} \mid \mathbb{F}_{\mathrm{PRED}\lambda\rightarrow_0} \supset \mathbb{F}_{\mathrm{PRED}\lambda\rightarrow_0} \mid (\forall x^{\mathbf{l}})\mathbb{F}_{\mathrm{PRED}\lambda\rightarrow_0}.$$

(v) *Postulates of* PRED$\lambda\!\rightarrow_0$

$$\begin{aligned}
\varphi\in\Gamma \quad &\Rightarrow \quad \Gamma \vdash \varphi;\\
\Gamma \vdash \varphi \supset \psi,\ \ \Gamma \vdash \varphi \quad &\Rightarrow \quad \Gamma \vdash \psi;\\
\Gamma, \varphi \vdash \psi \quad &\Rightarrow \quad \Gamma \vdash \varphi \supset \psi;\\
\Gamma \vdash \forall x^{\mathbf{l}}\varphi,\ \ t\in\mathbb{T}^{\mathbf{l}} \quad &\Rightarrow \quad \Gamma \vdash \varphi[x^{\mathbf{l}} := t];\\
\Gamma \vdash \varphi,\ \ x^{\mathbf{l}}\notin \mathrm{FV}(\Gamma) \quad &\Rightarrow \quad \Gamma \vdash \forall x^{\mathbf{l}}\varphi.
\end{aligned}$$

4.2. Definition. The mappings $[-]^i_r$ and Γ in B2.12 from $\mathbb{F}_{\mathrm{PRED}}$ into $\Lambda^s(\mathbf{\Xi}, \mathbf{L})$ and illative contexts respectively are extended to mappings from $\mathbb{F}_{\mathrm{PRED}\lambda\rightarrow_0}$ by altering some clauses and adding some clauses in the following way.

First we define $[\mathbf{A}_j]^i_r = A_j$ $(j = 1, 2)$, $[\mathbf{l}_1{\rightarrow}\mathbf{l}_2]^i_r = \mathbf{F}[\mathbf{l}_1]^i_r[\mathbf{l}_2]^i_r$. We will denote $[\mathbf{l}]^i_r$ by l. Then we define

$[x^{\mathbf{l}}]^i_r = x^l$, $\quad [\mathbf{t}_1\mathbf{t}_2]^i_r = [\mathbf{t}_1]^i_r[\mathbf{t}_2]^i_r$, $\quad [\forall x^{\mathbf{l}}\varphi]^1_r = \mathbf{\Xi}l(\lambda x^l.[\varphi]^1_r), [\forall x^{\mathbf{l}}\varphi]^2_r = \mathbf{G}l(\lambda x^l.[\varphi]^2_r)$,
$\Gamma(x^{\mathbf{l}}) = lx^l$, $\quad \Gamma(\forall x^{\mathbf{l}}\varphi) = \Gamma(\varphi) - \{lx^l\}$, $\quad \Gamma(\mathbf{t}_1\mathbf{t}_2) = \Gamma(\mathbf{t}_1) \cup \Gamma(\mathbf{t}_2)$,
$\Gamma^1_{r,s} =< \mathbf{L}A_1, \mathbf{L}A_2, \mathbf{F}A_1A_1f, \mathbf{F}A_1(\mathbf{F}A_2A_1)g, \mathbf{F}A_1\mathbf{H}(r{\circ}P), A_1a >$,
$\Gamma^2_{r,s} =< \mathbf{L}A_1, \mathbf{L}A_2, \mathbf{F}A_1A_1f, \mathbf{F}A_1(\mathbf{F}A_2A_1)g, \mathbf{F}A_1\mathbf{L}(r{\circ}P), A_1a >$,
$\Gamma^{i,+}_{r,s} = \Gamma^i_{r,s} \cup \{lx^l_1 \mid l \neq A_1\}$,
$[-]^i, \Gamma^i_s$ and $\Gamma^{i,+}_s$ are obtained from $[-]^i_r, \Gamma^i_{r,s}$ and $\Gamma^{i,+}_{r,s}$ by everywhere omitting r i.e. by replacing $r{\circ}P$ by P.

4.3. Proposition. $\mathcal{I}\mathbf{G}$ *is not complete for* PRED$\lambda\!\rightarrow_0$.

Proof. Let $\varphi = \forall x_1^{\mathbf{A}_1\rightarrow\mathbf{A}_1}\forall x_2^{\mathbf{A}_1}P(x_1x_2) \supset \forall x_2^{\mathbf{A}_1}Px_2$. We will show

$$\forall r[\Gamma^{2,+}_{r,s}, \Gamma(\varphi) \vdash_{\mathcal{I}\mathrm{G}} [\varphi]^2_r M \text{ some } M], \tag{$*$}$$

$$\forall_{\mathrm{PRED}\lambda\rightarrow_0}\varphi. \tag{$**$}$$

Proof of $(*)$. Let $X_1 = \mathbf{G}(\mathbf{F}A_1A_1)(\lambda x_1.\mathbf{G}A_1(\lambda x_2.(r{\circ}P)(x_1x_2)))$,
$Y_1 = \mathbf{G}A_1(\lambda x_2.(r{\circ}P)x_2), M = \lambda x^{X_1}.x^{X_1}I$ where $I = \lambda x^{A_1}.x^{A_1}$. We will prove $(*)$ by showing $\forall r[\Gamma^{2,+}_{r,s} \vdash_{\mathcal{I}\mathrm{G}} \mathbf{F}X_1Y_1M]$.

Note that the rules $\mathsf{G_e}$ and $\mathsf{G_i}$ imply the rules $\mathsf{F_e}$ and $\mathsf{F_i}$. Let r be a closed term in $\Lambda^s(\mathbf{\Xi}, \mathbf{L})$. We have by rule $\mathsf{F_i}$

$$\mathbf{L}A_1 \vdash \mathbf{F}A_1A_1I$$

and from this we get by rule $\mathsf{G_e}$ (taking
$X = \mathbf{F}A_1A_1, Y = \lambda x_1.\mathbf{G}A_1(\lambda x_2.(r{\circ}P)(x_1x_2)), Z = x(= x^{X_1}), V = I$)

$$\mathbf{L}A_1, X_1x \vdash Y_1(xI).$$

So we have

$$\mathbf{L}A_1, X_1x \vdash Y_1(Mx). \tag{1}$$

Moreover rule $\mathsf{F_e}$ yields

$$\mathbf{L}A_1, \mathbf{F}A_1A_1x_1, A_1x_2 \vdash A_1(x_1x_2).$$

Hence we get by applying rule $\mathsf{F_e}$ once more

$$\mathbf{L}A_1, \mathbf{F}A_1\mathbf{L}(r{\circ}P), \mathbf{F}A_1A_1x_1, A_1x_2 \vdash \mathbf{L}((r{\circ}P)(x_1x_2))$$

and by rule $\mathsf{G_L}$ twice

$$\mathbf{L}A_1, \mathbf{F}A_1\mathbf{L}(r{\circ}P) \vdash \mathbf{L}X_1. \tag{2}$$

From (1) and (2) we get by rule $\mathsf{F_i}$

$$\mathbf{L}A_1, \mathbf{F}A_1\mathbf{L}(r{\circ}P) \vdash \mathbf{F}X_1Y_1M.$$

This proves $(*)$. But $\not\vdash_{\mathrm{PRED}\lambda\to_0} \varphi$ because PRED$\lambda\to_0$ has no identity function in $\mathbb{T}^{\mathbf{F}\mathbf{A}_1\mathbf{A}_1}$. □

By similar examples one shows the need for other combinators or $\lambda-$terms. In fact all of these are needed in PRED$\lambda\to_0$ before completeness can be proved.

4.4. Definition (Predicate Calculus with Higher Type Functions, PRED$\lambda\to$).
Types and variables are as for PRED$\lambda\to_0$.
Terms of PRED$\lambda\to$

$$\begin{aligned}
\mathbb{T}^{\mathbf{l}} &= \mathbf{a} \mid \mathbb{V}^{\mathbf{l}} \mid \mathbb{T}^{\mathbf{l}_1\to\mathbf{l}}\mathbb{T}^{\mathbf{l}_1}, && \text{if } \mathbf{l} = \mathbf{A}_1;\\
\mathbb{T}^{\mathbf{l}} &= \quad \mathbb{V}^{\mathbf{l}} \mid \mathbb{T}^{\mathbf{l}_1\to\mathbf{l}}\mathbb{T}^{\mathbf{l}_1}, && \text{if } \mathbf{l} = \mathbf{A}_2;\\
\mathbb{T}^{\mathbf{l}} &= \mathbf{f} \mid \mathbb{V}^{\mathbf{l}} \mid \mathbb{T}^{\mathbf{l}_1\to\mathbf{l}}\mathbb{T}^{\mathbf{l}_1} \mid \lambda x^{\mathbf{A}_1}.\mathbb{T}^{\mathbf{A}_1}, && \text{if } \mathbf{l} = \mathbf{A}_1{\to}\mathbf{A}_1;\\
\mathbb{T}^{\mathbf{l}} &= \mathbf{g} \mid \mathbb{V}^{\mathbf{l}} \mid \mathbb{T}^{\mathbf{l}_1\to\mathbf{l}}\mathbb{T}^{\mathbf{l}_1} \mid \lambda x^{\mathbf{A}_1}\mathbb{T}^{\mathbf{A}_2\to\mathbf{A}_1}, && \text{if } \mathbf{l} = \mathbf{A}_1{\to}(\mathbf{A}_2{\to}\mathbf{A}_1);,\\
\mathbb{T}^{\mathbf{l}} &= \quad \mathbb{V}^{\mathbf{l}} \mid \mathbb{T}^{\mathbf{l}_1\to\mathbf{l}}\mathbb{T}^{\mathbf{l}_1} \mid \lambda x^{\mathbf{l}_2}.\mathbb{T}^{\mathbf{l}_3}, && \text{if } \mathbf{l} = \mathbf{l}_2{\to}\mathbf{l}_3, \mathbf{l}{\notin}\{\mathbf{A}_1{\to}\mathbf{A}_1, \mathbf{A}_1{\to}(\mathbf{A}_2{\to}\mathbf{A}_1)\}.
\end{aligned}$$

The well formed formulas of PRED$\lambda\to$ are defined as for PRED$\lambda\to_0$, but with the altered terms.
The postulates of PRED$\lambda\to$ are as for PRED$\lambda\to_0$, plus

$$\Gamma \vdash \varphi, \quad \varphi =_{\beta\eta} \psi \quad \Rightarrow \quad \Gamma \vdash \psi.$$

4.5. Remark. Ordinary predicate logic PRED can be expressed by the pure type system (PTS), see Barendregt[1992],

$\mathcal{S}$	$*^p, *^s, *^f, \Box^p, \Box^s$
$\mathcal{A}$	$*^p : \Box^p, *^s : \Box^s$
$\mathcal{R}$	$(*^p, *^p), (*^s, *^p), (*^s, \Box^p)$
	$(*^s, *^s, *^f), (*^s, *^f, *^f)$

The point is that $*^f$ is a sort for function spaces like $\mathbf{A}_1 \to \mathbf{A}_2$ or $\mathbf{A}_1 \to (\mathbf{A}_2 \to \mathbf{A}_2)$, with $\mathbf{A}_1, \mathbf{A}_2 \in *^s$, but not for higher function spaces like $(\mathbf{A}_1 \to \mathbf{A}_2) \to \mathbf{A}_2$. The system PRED$\lambda\to$ has higher function spaces and moreover it allows quantification over them. PRED$\lambda\to$ can be described more simply as the following PTS.

PRED$\lambda\to$

$\mathcal{S}$	$*^p, *^s, \Box^p, \Box^s$
$\mathcal{A}$	$*^p : \Box^p, *^s : \Box^s$
$\mathcal{R}$	$(*^p, *^p), (*^s, *^p), (*^s, \Box^p), (*^s, *^s)$

4.6. Definition. We extend the mappings $[-]^i_r$ and Γ of Definition 4.2 by

$$[\lambda x^{\mathbf{l}_1}.\mathbf{t}^{\mathbf{l}_2}]^i_r = \lambda x^{l_1}.[\mathbf{t}^{\mathbf{l}_2}]^i_r, \quad \Gamma(\lambda x^{\mathbf{l}_1}.\mathbf{t}^{\mathbf{l}_2}) = \Gamma(\mathbf{t}^{\mathbf{l}_2}) - \{l_1 x^{l_1}\}.$$

4.7. Proposition (Soundness of the interpretations for PRED$\lambda\to$). *Let $\Delta \cup \{\varphi\} \subseteq \mathbb{F}_{\mathrm{PRED}\lambda\to}$, then the following hold for all closed r.*

(i) $\Delta \vdash_{\mathrm{PRED}\lambda\to} \varphi \;\Rightarrow\; \Gamma^{1,+}_{r,s}, [\Delta]^1_r, \Gamma(\Delta, \varphi) \vdash_{\mathcal{I}\Xi} [\varphi]^1_r$.

(ii) $\Delta \vdash_{\mathrm{PRED}\lambda\to} \varphi \;\Rightarrow\; \Gamma^{2,+}_{r,s}, [\Delta]^2_r, \Gamma(\Delta, \varphi) \vdash_{\mathcal{I}\mathrm{G}} [\varphi]^2_r M$ *for some* M.

Proof. We shall not give the proof here. It is similar to the proof of Proposition B2.14 but longer because many more trivial cases have to be considered. □

Now in 4.8-4.12 we shall prove completeness of $\mathcal{I}\mathsf{G}$ for PRED$\lambda\to$ under the indirect translation $[-]^2$. (Completeness of $\mathcal{I}\Xi$ for PRED$\lambda\to$ under the direct translation $[-]^1$ also holds, but we shall not give the proof here.)

First we define in 4.8 an invariant $\mathcal{O}_3$, similar to $\mathcal{O}$ in B3.2 and $\mathcal{O}_1$ in 2.21. We have $\mathcal{O}_3 \subseteq \overline{\mathcal{O}_G}$, hence Lemma 2.18 yields that in a proof of $\Gamma \vdash_{\mathcal{I}\mathrm{G}} M$ where $\Gamma \subseteq \overline{\mathcal{O}_3}$ all cut formulas can be eliminated. Then the proof of completeness of $\mathcal{I}\mathsf{G}$ for PRED$\lambda\to$ is similar to the proof of completeness of $\mathcal{I}\mathsf{G}$ for PRED in 2.22 - 2.25.

4.8. Definition (Invariant $\mathcal{O}_3$ for $\mathcal{I}\mathsf{G}$).

$\mathcal{L}_3 = A_1 \mid A_2 \mid \mathsf{F}\mathcal{L}_3\mathcal{L}_3$;

$\mathcal{Q}_3 = \mathcal{L}_3 \mid \mathcal{P}_3$;

$\mathcal{V}_3^l = \{x_1^l, x_2^l, \ldots\}, \quad x_i^{l_1} \neq x_i^{l_2}$ if $l_1 \not\equiv l_2$;

$\mathcal{T}_3^l = a \mid \mathcal{V}_3^l \mid \mathcal{T}_3^{\mathsf{F}l_1 l}\mathcal{T}_3^{l_1}$, if $l \equiv A_1$;
$\mathcal{T}_3^l = \quad \mathcal{V}_3^l \mid \mathcal{T}_3^{\mathsf{F}l_1 l}\mathcal{T}_3^{l_1}$, if $l \equiv A_2$;
$\mathcal{T}_3^l = f \mid \mathcal{V}_3^l \mid \mathcal{T}_3^{\mathsf{F}l_1 l}\mathcal{T}_3^{l_1} \mid \lambda x^{A_1}.\mathcal{T}_3^{A_1}$, if $l \equiv \mathsf{F}A_1A_1$;
$\mathcal{T}_3^l = g \mid \mathcal{V}_3^l \mid \mathcal{T}_3^{\mathsf{F}l_1 l}\mathcal{T}_3^{l_1} \mid \lambda x^{A_1}.\mathcal{T}_3^{\mathsf{F}A_2A_1}$, if $l \equiv \mathsf{F}A_1(\mathsf{F}A_2A_1)$;
$\mathcal{T}_3^l = \quad \mathcal{V}_3^l \mid \mathcal{T}_3^{\mathsf{F}l_1 l}\mathcal{T}_3^{l_1} \mid \lambda x^{l_2}.\mathcal{T}_3^{l_3}$, if $l \equiv \mathsf{F}l_2l_3, l \notin \{\mathsf{F}A_1A_1, \mathsf{F}A_1(\mathsf{F}A_2A_1)\}$.

$\mathcal{P}_3 \equiv P\mathcal{T}_3^{A_1} \mid \mathsf{G}q(\lambda x^q.\mathcal{P}_3)$;
$\mathcal{H}_3 \equiv (Pt^{A_1})t^{Pt^{A_1}} \mid \Xi q(\lambda x^q.\mathcal{H}_3)$;
$\mathcal{G}_3 \equiv \mathsf{L}\mathcal{Q}_3 \mid A_i\mathcal{T}_3^{A_i} \mid \Xi q(\lambda x^q.\mathcal{G}_3)$;
$\mathcal{O}_3 \equiv \mathcal{H}_3 \mid \mathcal{G}_3$.

$\mathcal{V}_3^p = \{x_1^p, x_2^p, \ldots\} \quad , x_i^p = x_i^{sk(p)}, x_i^{p_1} \neq x_i^{p_2}$ if $sk(p_1) \not\equiv sk(p_2)$;

$sk(Pt^{A_1}) \equiv Px^{A_1}$ (i.e. $sk(Pt^{A_1})$ is independent of t^{A_1});
$sk(\mathsf{G}l(\lambda x^l.p)) \equiv \mathsf{G}l(\lambda x^l.sk(p))$;

$\mathcal{T}_3^p \equiv \mathcal{V}_3^p \mid \mathcal{T}_3^{\mathsf{G}l(\lambda x^l.p)}\mathcal{T}_3^l$ if $p \equiv Pt^{A_1}$;
$\mathcal{T}_3^p \equiv \mathcal{V}_3^p \mid \mathcal{T}_3^{\mathsf{G}l(\lambda x^l.p)}\mathcal{T}_3^l \mid \lambda x^q.\mathcal{T}_3^{p_1}$, if $p \equiv \mathsf{G}q(\lambda x^q.p_1))$.

4.9. Remarks. (i) Similar to Lemma 2.4(v) we now have $\mathcal{H}_3 = \mathcal{P}_3\mathcal{T}_3^{\mathcal{P}_3}$.

(ii) If $q \equiv l$ in $\mathsf{G}q(\lambda x^q.p_1)$ in the definition of $\mathcal{P}_3$, then x^q may occur free in p_1; but if $q \equiv p$ then $x^q \notin \mathrm{FV}(p_1)$. So $\mathsf{G}q(\lambda x^q.p_1)$ may be split into $\mathsf{G}l(\lambda x^l.p_1)$ and $\mathsf{G}p(\mathsf{K}p_1)$ and similarly for $\Xi q(\lambda x^q.g)$ in the definition of $\mathcal{G}_3$.

Lemma's 2.6 - 2.9 mutatis mutandis also hold for $\mathcal{O}_3$. (Replace l by q.) Moreover lemma's corresponding to Lemma's B3.7 - B3.9 hold. Replace $[-]^1$ by $[-]^2$ and in the lemma corresponding to B3.7 take $\mathcal{H}_3$ instead of $\mathcal{P}_3$.

4.10. Lemma. *Let $p, p' \in \mathcal{P}_3$. Then $pM = p'M' \;\Rightarrow\; p = p'$.*

Proof. Induction on p. □

4.11. Lemma. *Let $\Gamma \subseteq \overline{\mathcal{O}_3}$ and let $\Gamma \vdash_{\mathcal{IG}} N$ be a cut free proof. Then*

(i) *If the last applied rule in the proof is not the $\Xi_{\mathrm{i}}-$rule, nor the $\mathsf{G}_{\mathsf{L}}-$rule (modulo the $\beta\eta-$rule), then $N \in \overline{\mathcal{O}_3}$.*

(ii) *If $N = qt$ with $q \in \mathcal{Q}_3, t \in \mathcal{T}_G^q$ then $N \in \overline{\mathcal{O}_3}$.*

Proof. Similar to the proof of 2.24. □

4.12. Proposition (Completeness of $\mathcal{I}\mathbf{G}$ for PRED$\lambda\!\rightarrow$ relative to $[-]^2$).

$$\Gamma_s^{2,+}, [\Delta]^2, \Gamma(\Delta,\varphi) \vdash_{\mathcal{I}\mathrm{G}} [\varphi]^2 M \;\Rightarrow\; \Delta \vdash_{\mathrm{PRED}\lambda\rightarrow} \varphi.$$

Proof. We have $\mathcal{O}_3 \subseteq \overline{\mathcal{O}_G}$, hence by Lemma 2.18 we may assume that the proof of $\Gamma_s^{2,+}, [\Delta]^2, \Gamma(\Delta,\varphi) \vdash_{\mathcal{I}\mathrm{G}} [\varphi]^2 M$ is cut free. Then the proof is similar to the proof of completeness of $\mathcal{I}\mathbf{G}$ for PRED in Proposition 2.25. □

4.13. Remark (Higher Order Predicate Calculus). Let us define PRED$\lambda\omega$ as PRED$\lambda\!\rightarrow$ but with

$$\mathbb{L}_{\mathrm{PRED}\lambda\omega} = \mathbf{A}_1 \mid \mathbf{A}_2 \mid \mathbf{H} \mid \mathbb{L}_{\mathrm{PRED}\lambda\omega}\!\rightarrow\!\mathbb{L}_{\mathrm{PRED}\lambda\omega}.$$

Thus PRED$\lambda\omega$ allows quantification over all functions, but also over all predicates and propositions. As a PTS the system PRED$\lambda\omega$ is PRED$\lambda\!\rightarrow$ plus $(\Box^p, *^p), (\Box^p, \Box^p)$. Given an extension of $\Gamma^i_{r,s}$ to include $\vdash \mathbf{LH}$, we would expect to prove the completeness of $\mathcal{I}\mathbf{G}$ over PRED$\lambda\omega$, but we have not yet succeeded. In particular we have not been able to prove cut elimination for $\mathbf{LH}\!\in\!\Gamma^i_{r,s}$, so the methods employed here are not all available.

5. Conclusions and remarks

In BBD[1993] and in this paper we proved soundness and completeness for the direct and indirect interpretations of propositional and predicate calculus into illative systems:

(i) $\Delta \vdash_{\mathrm{PROP}} \varphi \;\Leftrightarrow\; \forall r[\ldots_1, [\Delta]^1_r \vdash_{\mathcal{I}\mathrm{P}} [\varphi]^1_r]$;
(ii) $\Delta \vdash_{\mathrm{PRED}} \varphi \;\Leftrightarrow\; \forall r[\ldots_1, [\Delta]^1_r \vdash_{\mathcal{I}\Xi} [\varphi]^1_r]$;
(iii) $\Delta \vdash_{\mathrm{PROP}} \varphi \;\Leftrightarrow\; \forall r \exists M[\ldots_2, [\Delta]^2_r \vdash_{\mathcal{I}\mathrm{F}} [\varphi]^2_r M]$;
(iv) $\Delta \vdash_{\mathrm{PRED}} \varphi \;\Leftrightarrow\; \forall r \exists M[\ldots_2, [\Delta]^2_r \vdash_{\mathcal{I}\mathrm{G}} [\varphi]^2_r M]$;

where $\ldots_i$ stands for $\Gamma^i_r(\Delta,\varphi)$ in the cases of PROP and for $\Gamma^{i,+}_{r,s}, \Gamma(\Delta,\varphi)$ in the cases of PRED.

5.1. Remark. The proof of soundness ($\Rightarrow$) was always simple. Completeness ($\Leftarrow$) was proved each time by specialising to $r = \mathbf{I}$. This is a very natural choice, but one may wonder if the proofs can be given for other r, especially for $r = \mathbf{K}$. Inspecting the 4 proofs for $r = \mathbf{I}$ in this paper and in BBD[1993] one can verify that all these proofs need only minor changes to yield proofs for $r = \mathbf{K}$.

Let us consider as an example the proof of completeness of $\mathcal{I}\mathbf{G}$ for PRED ((ii)$\Leftarrow$) taking $r = \mathbf{K}$. Each Pt^{A_1} in section 2 must be replaced by $\mathbf{K}(Pt^{A_1})$. (And a resulting $\mathbf{K}(Pt^{A_1})M$ we rewrite Pt^{A_1} again.) Let us denote for the moment the resulting set $\mathcal{P}_G$ by $\mathcal{P}_{G\mathbf{K}}$ and the set $\mathcal{P}_G$ of section 2 for $r = \mathbf{I}$ by $\mathcal{P}_{G\mathbf{I}}$. Then

$$\mathcal{P}_{G\mathbf{K}} = \mathbf{K}\mathcal{P}_{G\mathbf{I}}.$$

In Definition 2.2 we define now

$$\mathcal{T}^l_G \equiv \Lambda^s(\mathbf{\Xi}, \mathbf{L}) \text{ if } l \equiv \mathbf{K}(Pt^{A_1}).$$

As a result then $\mathcal{T}^p_G = \Lambda^s(\mathbf{\Xi}, \mathbf{L})$ for each $p \in \mathcal{P}_{G\mathbf{K}}$. These are nearly all changes that are needed to get a proof for $r = \mathbf{K}$ from the proof for $r = \mathbf{I}$ in section 2.

In an earlier version of this paper we were not yet able to give completeness proofs of $\mathcal{I}\mathbf{G}$ for PRED and $\mathcal{I}\mathbf{F}$ for PROP ((iv)$\Leftarrow$ and (iii)$\Leftarrow$) by specialising to $r = \mathbf{I}$. We then gave completeness proofs by taking $r = \mathbf{K}$, in a way different from the method sketched above. Let us consider now the earlier proof of completeness of $\mathcal{I}\mathbf{G}$ for PRED taking $r = \mathbf{K}$. From

$$[\varphi]^2_{\mathbf{K}\circ r} = \mathbf{K}[\varphi]^1_r$$

in Lemma B2.15 we got

$$[\varphi]^2_{\mathbf{K}} M = [\varphi]^1_{\mathbf{I}}. \qquad (*)$$

Moreover we had a conservativity result

$$\Gamma \vdash_{\mathcal{I}\mathrm{G}} M \;\Rightarrow\; \Gamma \vdash_{\mathcal{I}\Xi} M \text{ for 'suitable' } \Gamma \text{ and } M. \qquad (**).$$

Now the completeness proof was as follows

$$\begin{array}{lll}
\Gamma^{2,+}_{\mathbf{K},s}, [\Delta]^2_{\mathbf{K}}, \Gamma(\Delta, \varphi) \vdash_{\mathcal{I}\mathrm{G}} [\varphi]^2_{\mathbf{K}} M & \Rightarrow & (\text{by } *) \\
\Gamma^{1,+}_{\mathbf{I},s}, [\Delta]^1_{\mathbf{I}}, \Gamma(\Delta, \varphi) \vdash_{\mathcal{I}\mathrm{G}} [\varphi]^1_{\mathbf{I}} & \Rightarrow & (\text{by } **) \\
\Gamma^{1,+}_{\mathbf{I},s}, [\Delta]^1_{\mathbf{I}}, \Gamma(\Delta, \varphi) \vdash_{\mathcal{I}\Xi} [\varphi]^1_{\mathbf{I}} & \Rightarrow & (\text{by Proposition B3.14}) \\
\Delta \vdash_{\mathrm{PRED}} \varphi. & &
\end{array}$$

(To prove the conservativity result we had to show, in a way similar to the proof in the present paper, that the proof of $\Gamma \vdash_{\mathcal{I}\mathrm{G}} M$ could be made cut free.)

Also completeness of $\mathcal{I}\mathbf{F}$ relative to PROP for $[-]^2_{\mathbf{K}}$ followed from completeness of $\mathcal{I}\mathbf{P}$ relative to PROP for $[-]^1_{\mathbf{I}}$ by a conservativity result for $\mathcal{I}\mathbf{F}$ relative to $\mathcal{I}\mathbf{P}$.

We also proved completeness of $\mathcal{I}\mathbf{G}$ relative to PRED$\lambda\!\!\rightarrow$ for$[-]^2_{\mathbf{K}}$. Then completeness of $\mathcal{I}\mathbf{\Xi}$ relative to PRED$\lambda\!\!\rightarrow$ for$[-]^1_{\mathbf{I}}$ *followed* because system $\mathcal{I}\mathbf{G}$ is stronger then system $\mathcal{I}\mathbf{\Xi}$. That last completeness is not proved in the present paper.

5.2. Remark. We defined $\mathbf{P}, \mathbf{F}, \mathbf{G}$ in B2.3 in terms of the constant $\mathbf{\Xi}$ and in B2.7 we obtained the following result.
The systems $\mathcal{I}\mathbf{P}$, $\mathcal{I}\mathbf{\Xi}$, $\mathcal{I}\mathbf{F}$ and $\mathcal{I}\mathbf{G}$ are related as follows



where $\longrightarrow$ denotes nondecreasing strength, i.e. $s_1 \longrightarrow s_2$ means that for all Γ, X

$$\Gamma \vdash_{s_1} X \;\Rightarrow\; \Gamma \vdash_{s_2} X.$$

Moreover we had conservativity results like

$$\Gamma \vdash_{\mathcal{I}\mathsf{G}} M \;\Rightarrow\; \Gamma \vdash_{\mathcal{I}\Xi} M \text{ for 'suitable' } \Gamma \text{ and } M$$

mentioned above. But in the proofs of soundness and completeness in the present paper we did not use these results. So one may wonder if soundness and completeness remain valid if we define P, F and G to be constants. This is indeed the case (only the bases $\Gamma^1_{r,s}$ and $\Gamma^2_{r,s}$ must be rewritten). In the soundness proofs we did not use the fact that P, F and G were defined in terms of Ξ. Also completeness remains valid because the systems with P, F and G constant are less strong than the systems we considered. For example for system $\mathcal{I}\mathsf{F}$ then one does not have the relation $\mathsf{F} =_{\beta\eta} \lambda xyz.\Xi x(y \circ z)$.

5.3. Remark. In B4.1 we stated that it is possible to work with variants of the systems $\mathcal{I}\mathsf{P}$, $\mathcal{I}\Xi$, $\mathcal{I}\mathsf{F}$ and $\mathcal{I}\mathsf{G}$ based on β-conversion only. This still holds. One may think that in the proof of 2.9 (and the proof of 4.11) the $\eta-$axiom is used. This however is not the case, because by Definition 2.2 not only $ft^{A_1} \in \mathcal{T}_G^{A_1}$, but also $f \in \mathcal{T}_G^{\mathsf{F}A_1A_1}$ and similarly $g \in \mathcal{T}_G^{\mathsf{F}A_1(\mathsf{F}A_2A_1)}$.

References

Barendregt H., Bunder M. and Dekkers W.

[1993] Systems of illative combinatory logic complete for first-order propositional and predicate calculus, *Journal of Symbolic Logic* **58**, 769-788.

Barendregt H.

[1992] Lambda Calculi with Types, *Handbook of Logic in Computer Science, Vol II,* S. Abramski, D.M. Gabbay and T.S.E. Maibaum (eds.), Oxford University Press.