# THEORETICAL PEARLS

## *Applications of Plotkin-terms: partitions and morphisms for closed terms*

RICHARD STATMAN

*Department of Mathematics, Carnegie-Mellon University, Pittsburgh, Pennsylvania 15213, USA.*
(*e-mail:* Rick.Statman@andrew.cmu.edu)

HENK BARENDREGT

*Department of Computer Science, Catholic University, Box 9102, 6500 HC Nijmegen, The Netherlands.*
(*e-mail:* henk@cs.kun.nl)

**Abstract**

This theoretical pearl is about the closed term model of pure untyped lambda-terms modulo $\beta$-convertibility. A consequence of one of the results is that for arbitrary distinct combinators (closed lambda terms) $M, M', N, N'$ there is a combinator $H$ such that

$$HM = HM' \neq HN = HN'.$$

The general result, which comes from Statman [1998], is that uniformly r.e. partitions of the combinators, such that each "block" is closed under $\beta$-conversion, are of the form $\{H^{-1}\{M\}\}_{M \in \Lambda^\emptyset}$. This is proved by making use of the idea behind the so-called Plotkin-terms, originally devised to exhibit some global but non-uniform applicative behavior. For expository reasons we present the proof below. The following consequences are derived: a characterization of morphisms and a counter-example to the perpendicular lines lemma for $\beta$-conversion.

## 1. Introduction

We use notations from recursion theory and lambda calculus, see Rogers [1987] and Barendregt [1984].

NOTATION. (i) $\varphi_e$ is the $e$-th partial recursive function of one argument.
(ii) $W_e = \mathrm{dom}(\varphi_e) \subseteq \mathbb{N}$ is the r.e. set with index $e$.
(iii) $\Lambda$ is the set of lambda-terms and $\Lambda^\emptyset$ is the set of closed-lambda terms (combinators).
(iv) $\mathcal{W}_e = \{M \in \Lambda^\emptyset \mid \#M \in W_e\} \subseteq \Lambda^\emptyset$; here $\#M$ is the code of the term $M$.

1.1. DEFINITION. (i) Inspired by Visser [1980] we define a *Visser-partition* (V-partition) of $\Lambda^\emptyset$ to be a family $\{\mathcal{W}_e\}_{e \in S}$ such that

(1) $S \subseteq \mathbb{N}$ is an r.e. set
(2) $\forall e{\in}S\ \forall M, N\ (M{\in}\mathcal{W}_e\ \&\ N = M)\ \Rightarrow\ N{\in}\mathcal{W}_e$.
(3) $\mathcal{W}_e \cap \mathcal{W}_{e'} \neq \emptyset\ \Rightarrow\ \mathcal{W}_e = \mathcal{W}_{e'}$.

(ii) A family $\{\mathcal{W}_e\}_{e\in S}$ is a *pseudo*-V-partition if it satisfies just 1 and 2.

1.2. DEFINITION. Let $\{\mathcal{W}_e\}_{e\in S}$ be a V-partition.

1. The partition is said to be *covering* if $\bigcup_{e\in S} \mathcal{W}_e = \Lambda^\emptyset$.
2. The partition is said to be *inhabited* if $\forall e{\in}S\ \mathcal{W}_e \neq \emptyset$.
3. A V-partition $\{\mathcal{W}_e\}_{e\in S'}$ is said to be *(extensionally) equivalent* with $\{\mathcal{W}_e\}$ if these families define the same collection of non-empty sets, i.e. if

$$\{\mathcal{W}_e \mid e{\in}S\ \&\ \mathcal{W}_e \neq \emptyset\} = \{\mathcal{W}_e \mid e{\in}S'\ \&\ \mathcal{W}_e \neq \emptyset\}.$$

1.3. EXAMPLE. Let $H$ be some given combinator. Define

$$\mathcal{W}_{e(M,H)} = \{N{\in}\Lambda^\emptyset \mid HN = HM\},$$

Then $\{\mathcal{W}_e\}_{e\in S_H}$, with $S_H = \{e(M,H) \mid M{\in}\Lambda^\emptyset\}$, is an example of a covering and inhabited V-partition. We denote this V-partition by $\{\mathcal{W}_{e(M,H)}\}_{M\in\Lambda^\emptyset}$.

1.4. PROPOSITION. (i) *Every V-partition is effectively equivalent to an inhabited one.*

(ii) *Every V-partition can effectively be extended to a covering one.*

PROOF. (i) Given $\{\mathcal{W}_e\}_{e\in S}$ define $S' = \{e{\in}S \mid \mathcal{W}_e \neq \emptyset\}$. Then $\{\mathcal{W}_e\}_{e\in S'}$ is the required modified partition.

(ii) Given $\{\mathcal{W}_e\}_{e\in S}$ define

$$\mathcal{W}_{e(M)} = \{N \mid N = M\ \vee\ \exists e{\in}S\ M, N{\in}\mathcal{W}_e\}.$$

Then $\{\mathcal{W}_{e(M)}\}_{M\in\Lambda^\emptyset}$ is the required V-partition. ■

The main theorem comes in two version. The second more sharp version is needed for the construction of so called inevitably consistent equations, see Statman [1999].

1.5. THEOREM (Main theorem). (i) *Let $\{\mathcal{W}_e\}_{e\in S}$ be a V-partition. Then one can construct effectively a combinator $H$ such that for all $M, N{\in}\Lambda^\emptyset$*

$$HM = HN\ \Leftrightarrow\ M = N\ \vee\ \exists e{\in}S\ M, N{\in}\mathcal{W}_e. \qquad (*)$$

*The construction of $H$ is effective in the code of the underlying r.e. set $S$.*

(ii) *Let $\{\mathcal{W}_e\}_{e\in S}$ be a pseudo-V-partition. Then one can construct effectively a combinator $H$ such that if $\{\mathcal{W}_e\}_{e\in S}$ is an actual V-partition, then $(*)$ holds.*

The theorem will be proved in §2. It has several consequences. In order to state these we have to formulate the notion of morphism on $\Lambda^\emptyset$ and the so-called perpendicular lines lemma.

1.6. DEFINITION. Let $\varphi : \Lambda^\emptyset{\to}\Lambda^\emptyset$ be a map. Then $\varphi$ is a *morphism* if

1. $\varphi(M) = \mathsf{Ec}_{f(\#M)}$, for some recursive function $f$.
2. $M = N\ \Rightarrow\ \varphi(M) = \varphi(N)$.

1.7. LEMMA. (i) *Let $F$ be a combinator and define $\varphi_H(M) \equiv HM$. Then $\varphi_H$ is a morphism.*

(ii) *Let $F, G$ be combinators such that for all $M \in \Lambda^\emptyset$ there exists a unique $N \in \Lambda^\emptyset$ with $FM = GN$. Then there is a map $\varphi_{F,G}$ such that $FM = G\varphi_{F,G}(M)$, for all $M$, which is a morphism.*

PROOF. (i) For the coding $\#$ let app be the recursive function such that $\#(PQ) = \mathrm{app}(\#P, \#Q)$. Define $f(m) = \mathrm{app}(\#H, m)$. Then $\varphi_H(M) = \mathsf{Ec}_{f(\#M)}$. It is obvious that $\varphi_H$ preserves $\beta$-equality.

(ii) Let $R(m,n)$ be an r.e. relation. The we have $R(m,n) \Leftrightarrow \exists z\ T(m,n,z)$, for some recursive $T$. Let $< n, z >$ be a recursive pairing with recursive inverses $< n, z > .0 = n, < n, z > .1 = z$. Define ($\mu$ is the least number operator)

$$\iota_n.R(m,n) = (\mu p.T(m, p.0, p.1)).0.$$

Then $\exists n \in \mathbb{N} R(m,n) \Rightarrow R(m, \iota_n.R(m,n))$. In order to construct the morphism $\varphi_{F,G}$, define

$$f(m) = \iota_n.F(\mathsf{Ec}_m) = G(\mathsf{Ec}_n).$$

By the assumption (existence) $f$ is total. Define $\varphi_{F,G}(M) = \mathsf{Ec}_{f(\#M)}$. Now $f(\#M) = n \Rightarrow F(\mathsf{Ec}_n) = G(\mathsf{Ec}_n)$. Therefore $FM = G\varphi_{F,G}(M)$, for all $M$. The condition

$$M = M' \Rightarrow \varphi_{F,G}(M) = \varphi_{F,G}(M')$$

holds by the assumption (unicity). ■

One may wonder whether dropping the unicity condition in lemma 1.7 (ii) one may obtain a morphism by making a right uniformization. This is not the case.

1.8. PROPOSITION. *There exists combinators $F, G$ such that $\forall M \exists N\ FM = GN$ but without any morphism satisfying $\forall M\ FM = G\varphi(N)$.*

PROOF. Let $\Delta = \mathsf{Y}\Omega$ and define $F = \lambda x.\langle x, \Delta, \mathsf{I}\rangle$ and $G = \lambda y.\langle \mathsf{E}y, y\Omega\Delta, y\mathsf{I}\rangle$. Then, see Statman [1986],

$$FM =_\beta GN \Leftrightarrow (N =_\beta c_n \vee N =_\beta \mathsf{I}) \ \&\ \mathsf{E}N =_\beta M. \tag{1}$$

Any morphism $\varphi$ such that $FM = G\varphi(M)$ would solve the convertibility problem recursively: one has by (1)

$$M = M' \Leftrightarrow \varphi(M) = \varphi(M'), \tag{2}$$

and since $\varphi(M), \varphi(M')$ have nf's by (1), the RHS of (2) is decidable. ■

1.9. PROPOSITION. *Not every morphism is of the form $\varphi_H$.*

PROOF. Let $F, G \in \Lambda^\emptyset$ be such that $F \circ G = \mathsf{I}$. Then $F, G$ determine a so-called *inner model* $[\![\ ]\!] = [\![\ ]\!]^{F,G}$ as follows.

$$\begin{aligned} [\![x]\!] &= x; \\ [\![PQ]\!] &= F[\![P]\!][\![Q]\!]; \\ [\![\lambda x.P]\!] &= G(\lambda x.[\![P]\!]). \end{aligned}$$

Using the condition on $F, G$ it can be proved that

$$M =_\beta N \;\Rightarrow\; [\![M]\!] = [\![N]\!].$$

Therefore defining $\varphi(M) = [\![M]\!]$ we obtain a morphism.

Now take $F \equiv \lambda y.u\mathsf{I},\ \Gamma \equiv \lambda xy.yx$. Then indeed $F \circ G = \mathsf{I}$ and for the resulting inner model one has $[\![I]\!] = \lambda y.y\mathsf{I}$ and $[\![\Omega]\!] = (\lambda y.y(\lambda z.z\mathsf{I}z))\mathsf{I}(\lambda y.y(\lambda z.z\mathsf{I}z))$.

Suppose towards a contradiction that the resulting $\varphi$ is of the form $\varphi_H$. Then $H\mathsf{I} = \lambda y.y\mathsf{I}$, so $H$ is solvable and hence has a hnf $\lambda x_1 \ldots x_n.\cdot_i M_1 \ldots M_m$. But $H\Omega = (\lambda y.y(\lambda z.z\mathsf{I}z))\mathsf{I}(\lambda y.y(\lambda z.z\mathsf{I}z))$, which is unsolvable. Therefore the head-variable $x_i$ is $x_1$. But then $H\Omega = \lambda x_2 \ldots x_n.\Omega M_1^* \ldots M_m^*$ which is not of the correct form. ■

The following is a corollary to the main theorem.

1.10. Corollary. *Every morphism $\varphi$ is of the form $\varphi_{F,G}$.*

Proof. Let $\varphi$ be a given morphism. Define

$$\mathcal{W}_{e(N)} = \{Z \mid \exists M{\in}\Lambda^\emptyset\ [\varphi(M) = N \;\&\; [Z = \langle \mathbf{c}_0, M\rangle \;\vee\; Z = \langle \mathbf{c}_1, N\rangle]]\}.$$

Then $\{\mathcal{W}_{e(N)}\}$ is a V-partition. By the main theorem there exists an $H$ such that

$$\begin{aligned} H\langle \mathbf{c}_0, M\rangle = H\langle \mathbf{c}_1, N\rangle \quad &\Leftrightarrow \quad \langle \mathbf{c}_0, M\rangle = \langle \mathbf{c}_1, N\rangle \vee N = \varphi(M) \\ &\Leftrightarrow \quad N = \varphi(M). \end{aligned}$$

Define

$$\begin{aligned} F &= \lambda m.H\langle \mathbf{c}_0, m\rangle; \\ G &= \lambda n.H\langle \mathbf{c}_1, n\rangle. \end{aligned}$$

Then $FM = GN \;\Leftrightarrow\; N = \varphi(M)$. Therefore $\varphi = \varphi_{F,G}$. ■

Note that for a given morphism $\varphi$ one can define by

$$\mathcal{W}_{e(M,\varphi)} = \{N{\in}\Lambda^\emptyset \mid \varphi(M) = \varphi(N)\}.$$

This is an inhabited V-partition. It is not difficult to show that that each V-partition is equivalent to one of the form $\{\mathcal{W}_{e(M,\varphi)}\}$. Note that $\{\mathcal{W}_{e(M,H)}\} = \{\mathcal{W}_{e(M,\varphi_H)}\}$, see lemma 1.7. The following result shows that covering V-partitions are always of this more restricted form.

1.11. Corollary. *If $\{\mathcal{W}_e\}$ is a covering V-partition, then $\{\mathcal{W}_e\}$ is equivalent to $\{\mathcal{W}_{e(M,H)}\}_{M\in\Lambda^\emptyset}$ for some $H$, effectively found from $\{\mathcal{W}_e\}$.*

Proof. Let $H$ be the combinator constructed effectively from $\{\mathcal{W}_e\}$. We will show that $\mathcal{W}_{e(M,H)} = \{N \mid HN = HM\}$ is equivalent to $\{\mathcal{W}_e\}$. Claim. For $N{\in}\mathcal{W}_e$ one has $\mathcal{W}_e = \mathcal{W}_{e(M,H)}$. Indeed,

$$\begin{aligned} N{\in}\mathcal{W}_e \quad &\Leftrightarrow \quad M = N \;\vee\; M, N{\in}\mathcal{W}_e \\ &\Leftrightarrow \quad HN = HM \\ &\Leftrightarrow \quad N{\in}\mathcal{W}_{e(M,H)}. \end{aligned}$$

Therefore, noting that $M \in \mathcal{W}_{e(M,H)}$,

$$\{\mathcal{W}_e \mid M \in \Lambda^\emptyset, \mathcal{W}_e \neq \emptyset\} \subseteq \{\mathcal{W}_{e(M,H)} \mid \mathcal{W}_{e(M,H)} \neq \emptyset, M \in \Lambda^\emptyset\}.$$

The converse inclusion holds also, since every $M$ belongs to some $\mathcal{W}_e$ and hence $\mathcal{W}_{e(M,H)} = \mathcal{W}_e$ for this $e$. ■

The following theorem states that if a combinator, seen as function of $n$ arguments, is constant—modulo Böhm-tree equality—on $n$ perpendicular lines, then it is constant everywhere.

1.12. THEOREM (Perpendicular lines lemma). *Let $F$ be a combinator. Suppose that for $n \in \mathbb{N}$ there are combinators $M_{ij}$, $1 \leq i \neq j \leq n$, and $N_1, \ldots, N_n$ such that for all combinators $Z$ one has ($\cong$ denotes Böhm-tree equality, i.e. $M \cong N \Leftrightarrow BT(M) = BT(N)$)*

$$\begin{array}{cccccccc} F & Z & M_{12} & \ldots & M_{1n-1} & M_{1n} & \cong & N_1; \\ F & M_{21} & Z & \ldots & M_{2n-1} & M_{2n} & \cong & N_2; \\ & & & \ldots & & & & \\ & & & \ldots & & & & \\ F & M_{n1} & M_{n2} & \ldots & M_{nn-1} & Z & \cong & N_n. \end{array}$$

*Then for all $P_1, \ldots, P_n \in \Lambda^\emptyset$ one has*

$$FP_1 \ldots P_n \cong N_1 (\cong N_2 \cong \ldots \cong N_n).$$

PROOF. This is the restriction to closed terms of a theorem in Barendregt [1984], theorem 14.4.12, having the same proof. ■

1.13. COROLLARY. *The perpendicular lines lemma is false for any $n > 1$, if $\cong$ is replaced by $=_\beta$.*

PROOF (For $n = 1$ the perpendicular lines lemma is trivially true for $=_\beta$.). Let $n > 1$. For notational simplicity we assume $n = 2$ and give a counter example. Define

$$\begin{array}{lcl} \mathcal{W}_{e_1} & = & \{N \in \Lambda^\emptyset \mid N = \langle \mathsf{S}, \mathsf{S} \rangle\} \\ \mathcal{W}_{e_2} & = & \{N \in \Lambda^\emptyset \mid \exists Z \in \Lambda^\emptyset [N = \langle \mathsf{I}, Z \rangle \ \vee \ N = \langle Z, \mathsf{I} \rangle]\} \end{array}$$

Then $\{\mathcal{W}_e\}_{e \in \{e_1, e_2\}}$ is a V-partition. Let $H$ be the combinator obtained from this partition by the main theorem. Then for all $Z \in \Lambda^\emptyset$

$$H\langle \mathsf{S}, \mathsf{S} \rangle \neq H\langle \mathsf{I}, Z \rangle = H\langle Z, \mathsf{I} \rangle.$$

Now define $F \equiv \lambda xy.H\langle x, y \rangle$. Then for all $Z \in \Lambda^\emptyset$

$$F\,\mathsf{S}\,\mathsf{S} \neq F\,\mathsf{I}\,Z = F\,Z\,\mathsf{I}.$$

This is indeed a counterexample. ■

We do believe the conjecture in Barendregt [1984], stating that the perpendicular line lemma with $\cong$ replaced by $=_\beta$ is correct for open terms.

## 2. Proof of the main theorem

In order to prove the main theorem 1.5, let a V-partition determined by $S$ be fixed in this section. By proposition 1.4 it may be assumed that the partition is inhabited.

2.1. LEMMA. *Let $\{\mathcal{W}_e\}_{e\in S}$ be an inhabited V-partition.*

(i) *There exists a total recursive function $f = f_S$ such that*

$$\forall e{\in}S\ W_e = \{f((2e+1)2^n) \mid n{\in}\mathbb{N}\}.$$

(ii) *There exists a combinator $\mathsf{E}^S$ such that*

$$\forall e{\in}S\ \mathcal{W}_e = \{\mathsf{E}^S\mathbf{c}_{(2e+1)2^n} \mid n{\in}\mathbb{N}\}.$$

PROOF. (i) By elementary recursion theory there exists a recursive function $h$ such that $W_e = \mathrm{Range}(\varphi_{h(e)})$ and $\varphi_{h(e)}$ is total, for all $e{\in}S$. Observing that $e, n$ are uniquely determined by $k = (2e+1)2^n$, define $f$ by $f(0) = 0$, $f((2e+1)2^n) = \varphi_{h(e)}(n)$.

(ii) Take $\mathsf{E}^S = \mathsf{E} \circ F_S$, where $F_S$ lambda defines $f_S$ and $\mathsf{E}\mathbf{c}_{\#M} = M$ for all $M{\in}\Lambda^\emptyset$. ■

2.2. DEFINITION. (i) Define

$$\begin{aligned} \mathtt{odd}(0) &= 0; \\ \mathtt{odd}((2e+1)2^n) &= 2e+1. \end{aligned}$$

(ii) Define $M \sim N$ iff $M = N \ \vee\ M = \mathsf{E}_m, N = \mathsf{E}_n$ and $\mathtt{odd}(m) = \mathtt{odd}(n)$, for some $m, n$.

Notice that $M \sim N$ iff $M = N$ or $\exists e{\in}S M, N{\in}\mathcal{W}_e$. Therefore we have to prove that there exists a combinator $H$ such that

$$HM = HN \ \Leftrightarrow\ M \sim N.$$

The proof consists in constructing a combinator $H = H^S$ such that

1. $M \sim N \ \Rightarrow\ HM = HN$, proposition 2.4;
2. $HM = HN \ \Rightarrow\ M \sim N$, proposition 2.9.

The second part of the main theorem easily follows by inspecting the proof.

2.3. DEFINITION. (i) Define

$$\begin{aligned} T &\equiv \lambda xyz.xy(xyz); \\ A &\equiv \lambda fgxyz.fx(a(\mathsf{E}x))[f(\mathsf{S}^+x)y(g(\mathsf{S}^+x))z]; \\ B &\equiv \lambda fgx.f(\mathsf{S}x)(a(\mathsf{E}(Tx))(g(\mathsf{S}^+x))(gx). \end{aligned}$$

(ii) By the double fixed-point theorem there exists terms $F, G$ such that

$$\begin{aligned} F &\twoheadrightarrow AFG; \\ G &\twoheadrightarrow BFG. \end{aligned}$$

To be explicit, write

$$\begin{array}{lcl} D & \equiv & (\lambda xy.y(xxy)); \\ Y & \equiv & DD; \\ G & \equiv & Y(\lambda u.B(Y(\lambda v.Auv))u); \\ F & \equiv & Y(\lambda u.AuG). \end{array}$$

(iii) Finally define

$$H \equiv \lambda xa.F\mathbf{c}_1(ax)(G\mathbf{c}_1).$$

NOTATION. Write

$$\begin{array}{rcl} F_k & \equiv & F\mathbf{c}_k; \\ G_k & \equiv & G\mathbf{c}_k; \\ \mathsf{E}_k & \equiv & \mathsf{E}\mathbf{c}_k; \\ a_k & \equiv & a\mathsf{E}_k; \\ H_k[\ ] & \equiv & F_k[\ ]G_k; \\ C_k[\ ] & \equiv & F_k a_k([\ ]G_k). \end{array}$$

Note that by construction

$$\begin{array}{rcl} F_kMN & \twoheadrightarrow & F_k a_k(F_{k+1}MG_{k+1}N); \\ G_k & \twoheadrightarrow & F_{k+1}a_{2k}G_{k+1}G_k. \end{array}$$

By reducing $F$, respectively $G$, it follows that

$$\begin{array}{lcll} H_k[a_p] & \equiv & F_k a_p G_k \twoheadrightarrow C_k[H_{k+1}[a_p]] & (1) \\ H_k[a_k] & \equiv & F_k a_k G_k \twoheadrightarrow C_k[H_{k+1}[a_{2k}]] & (2) \end{array}$$

2.4. PROPOSITION. $M \sim N \Rightarrow HM = HN$.

PROOF. By lemma 2.1 it suffices to show $H\mathsf{E}_k = H\mathsf{E}_{2k}$ for all $k$.

$$\begin{array}{rcll} H\mathsf{E}_k & = & \lambda a.H_1[a_k] & \\ & = & \lambda a.C_1[C_2[\ldots C_{k-1}[H_k[a_k]]..]], & \text{by (1)}, \\ & = & \lambda a.C_1[C_2[\ldots C_{k-1}[C_k[H_k[a_{2k}]]]..]], & \text{by (2)}, \\ H\mathsf{E}_{2k} & = & \lambda a.H_1[a_{2k}] & \\ & = & \lambda a.C_1[C_2[\ldots C_{k-1}[C_k[H_k[a_{2k}]]]..]], & \text{by (1)}. \ \blacksquare \end{array}$$

As a piece of art we exhibit in more detail the reduction flow (contracted redexes are underlined).

$$\begin{array}{l}
\underline{H\mathsf{E}_k}\\
\lambda a.\underline{F_1 a_k G_1}\\
\lambda a.F_1 a_1(\underline{F_2 a_2 G_2} G_1)\\
\lambda a.F_1 a_1(F_2 a_2(\underline{F_3 a_k G_3} G_2)G_1)\\
\ldots\\
\lambda a.F_1 a_1(F_2 a_2(F_3 a_3(\ldots(F_k a_k G_k G_{k-1})\ldots)G_2)G_1) \equiv\\
\lambda a.F_1 a_1(F_2 a_2(F_3 a_3(\ldots(F_k a_k \quad \underline{G_k} \quad G_{k-1})\ldots)G_2)G_1)\\
\lambda a.F_1 a_1(F_2 a_2(F_3 a_3(\ldots(F_k a_k(F_{k+1} a_{2k} G_{k+1} G_k)G_{k-1})\ldots)G_2)G_1)
\end{array}$$

And also

$$\begin{array}{l}
H\mathsf{E}_{2k} \twoheadrightarrow \ldots \twoheadrightarrow\\
\lambda a.F_1 a_1(F_2 a_2(F_3 a_3(\ldots(F_k a_k(F_{k+1} a_{2k} G_{k+1} G_k)G_{k-1})\ldots)G_2)G_1)
\end{array}$$

For the converse implication we need the fine structure of the reduction.

2.5. Definition. Define

$$\begin{array}{lcl}
D_k^0[M] & \equiv & F_k(aM) \equiv Y(\lambda u.AuG)\mathbf{c}_k(aM)\\
D_k^1[M] & \equiv & (\lambda y.y(DDy))(\lambda u.AuG)\mathbf{c}_k(aM)\\
D_k^2[M] & \equiv & (\lambda u.AuG)F_k(aM)\\
D_k^3[M] & \equiv & AFG\mathbf{c}_k(aM)\\
D_k^4[M] & \equiv & (\lambda gxyz.F_x(a\mathsf{E}_x)(F_{\mathsf{S}^+ x}y(g(\mathsf{S}^+ x))z))G\mathbf{c}_k(aM)\\
D_k^5[M] & \equiv & (\lambda xyz.F_x(a\mathsf{E}_x)(F_{\mathsf{S}^+ x}yG_{\mathsf{S}^+ x}z))\mathbf{c}_k(aM)\\
D_k^6[M] & \equiv & (\lambda yz.F_k(a\mathsf{E}_k)(F_{\mathsf{S}^+ \mathbf{c}_k}yG_{\mathsf{S}^+ \mathbf{c}_k}z))(aM)\\
D_k^7[M] & \equiv & (\lambda z.F_k(a\mathsf{E}_k)(F_{\mathsf{S}^+ \mathbf{c}_k}(aM)G_{\mathsf{S}^+ \mathbf{c}_k}z))
\end{array}$$

2.6. Lemma. *Let $F_k(aM)N$ head-reduce in $8p+q$ steps to $W$. Then*

$$\begin{array}{lcll}
W & \equiv & D_k^q[M]N, & \text{if } p=0;\\
 & \equiv & D_k^q[\mathsf{E}_k]((H_{k+1}[\mathsf{E}_k])^{p-1}(H_{k+1}[M]N)), & \text{else.}
\end{array}$$

Proof. Note that $F_k(aM)N \equiv D_k^0[M]N$. Moreover,

$$\begin{array}{lcll}
D_k^q[M]N & \to_h & D_k^{q+1}[M]N, & \text{for } q<7;\\
D_k^7[M]N & \to_h & D_k^0[\mathsf{E}_k](H_{k+1}[M]N). &
\end{array}$$

The rest is clear. At steps 16, 24 we obtain for example

$$\begin{array}{rcl}
D_k^7[\mathsf{E}_k](H_{k+1}[M]N) & \to_h & D_k^0[\mathsf{E}_k]((H_{k+1}[\mathsf{E}_k])(H_{k+1}[M]G_k)).\\
D_k^7[\mathsf{E}_k]((H_{k+1}[\mathsf{E}_k])(H_{k+1}[M]G_k)) & \to_h & D_k^0[\mathsf{E}_k]((H_{k+1}[\mathsf{E}_k])^2(H_{k+1}[M]G_k)).\ \blacksquare
\end{array}$$

Remember that a standard reduction $\sigma{:}M \twoheadrightarrow_s N$ always consists of a head-reduction followed by an internal reduction:

$$\sigma{:}M \twoheadrightarrow_h W \twoheadrightarrow_i N.$$

NOTATION. Write $M =_{s\leq n} N$ if there are standard reductons of length $\leq n$ from $M$ respectively $N$ to a common reduct $Z$. Similarly $M =_{i\leq n} N$ for internal standard reductions. Also the notations $=_{s<n}$ and $=_{i<n}$ will be used.

2.7. LEMMA. (i) $D_k^q[M]N =_{i\leq n} D_k^{q'}[M']N' \;\Rightarrow\; q = q' \;\&\; N =_{s\leq n} N'$.
(ii) $D_k^q[M]N =_{i\leq n} D_k^q[M']N' \;\&\; q < 7 \;\Rightarrow\; M =_{s\leq n} M'$.
(iii) $D_k^7[M]N =_{i\leq n} D_k^7[M']N' \;\Rightarrow\; H_{k+1}[M] =_{s\leq n} H_{k+1}[M']$.

PROOF. (i) Suppose $D_k^q[M]N =_{i\leq n} D_k^{q'}[M']N'$. Then By observing where the free variable $a$ occurs one can conclude that $q = q'$. Since the reductions to a common reduct are internal, the positions of $N, N'$ are not changed and hence $N =_{s\leq n} N'$.
(ii) Obvious from the definition of $D_k^q$.
(iii) In this case it follows that

$$D_k^0[\mathsf{E}_k](H_{k+1}[M]z) =_{i\leq n} D_k^0[\mathsf{E}_k](H_{k+1}[M']z).$$

The conclusion $H_{k+1}[M] =_{s\leq n} H_{k+1}[M']$ depends on the fact that there are the free variables $z$ to mark the residuals. ■

2.8. LEMMA. *Suppose* $G_k =_{s\leq n} (H_{k+1}[\mathsf{E}_k])^d(H_{k+1}[M]G_k)$. *Then*

$$H_{k+1}[\mathsf{E}(T\mathbf{c}_k)] =_{s<n} H_{k+1}[M].$$

PROOF. By induction on $d$. If $d = 0$, then we have $G_k =_{s\leq n} H_{k+1}[M]G_k$. So there are standard reductions of these two terms to a common reduct. Observe that the head-reduction starting with $G_k$ begins as follows.

$$\begin{aligned}
G_k &\equiv Y(\lambda u.B(Y(\lambda v.Avu))u)\mathbf{c}_k\\
&\to_h (\lambda x.x(Yx))(\lambda u.B(Y(\lambda v.Avu))u)\mathbf{c}_k\\
&\to_h (\lambda u.B(Y(\lambda v.Avu))u)G\mathbf{c}_k\\
&\to_h BFG\mathbf{c}_k\\
&\to_h (\lambda gx.F(\mathsf{S}^+k)(a(\mathsf{E}^S(Tx)))(g(\mathsf{S}^+k))(gx)G\mathbf{c}_k\\
&\to_h (\lambda x.F(\mathsf{S}^+k)(a(\mathsf{E}^S(Tx)))(G(\mathsf{S}^+k))(Gx))\mathbf{c}_k\\
&\to_h F(\mathsf{S}^+k)(a(\mathsf{E}^S(T\mathbf{c}_k)))(G(\mathsf{S}^+k))(G\mathbf{c}_k).
\end{aligned}$$

The heads of these terms are not of order 0 except the last one. But $H_{k+1}[X]$ is always of order 0. Therefore the mentioned standard reduction of $G_k$ goes at least to this last term $H_{k+1}[\mathsf{E}^S(T\mathbf{c}_k)]G_k$. But then $H_{k+1}[\mathsf{E}^S(T\mathbf{c}_k)] =_{s<n} H_{k+1}[M]$.

If $d > 0$, then start the same argument as above, but at the intermediate conclusion

$$H_{k+1}[\mathsf{E}^S(T\mathbf{c}_k)]G_k =_{s<n} (H_{k+1}[\mathsf{E}_k])^d(H_{k+1}[M]G_k),$$

one preceeds by concluding that

$$G_k =_{s<n} H_{k+1}[\mathsf{E}_k]^{d-1}(H_{k+1}[M]G_k)$$

and uses the induction hypothesis. ■

2.9. PROPOSITION. $H_k[M] = H_k[N] \;\Rightarrow\; M \sim N$.

PROOF. By the standardization theorem it suffices to show for all $n$ that

$$\forall k{\in}\mathbb{N}[H_k[M] =_{s\leq n} H_k[N] \;\Rightarrow\; M \sim N].$$

This will be done by induction on $n$. From $H_k[M] =_{s\leq n} H_k[N]$ it follows that

$$\begin{array}{lllll} H_k[M] & \twoheadrightarrow_h & W_M & \twoheadrightarrow_i & Z \\ H_k[N] & \twoheadrightarrow_h & W_N & \twoheadrightarrow_i & Z. \end{array}$$

for some $W_M, W_N, Z$.

Case 1. $W_M, W_N$ are both reached after $< 8$ steps. Then by lemma 2.6 $W_M \equiv D_k^q[M]G_k, W_N \equiv D_k^{q'}[N]G_k$. By lemma 2.7(i) it follows that $q = q'$. If $q < 7$, then by 2.7(ii) one has $M = N$ so $M \sim N$. If $q = 7$, then by 2.7(iii) one has $H_{k+1}[M] =_{s<n} H_{k+1}[N]$ and by the induction hypothesis one has $M \sim N$.

Case 2. $W_M$ is reached after $p \geq 8$ steps and $W_N$ after $q < 8$ steps. Then $p = 8d + q$ and, keeping in mind lemma 2.7(i), it follows that $W_M \equiv D_k^q[M]G_k, W_N \equiv D_k^q[E_k]R$, $G_k =_{s<n} R$, where $R \equiv (H_{k+1}[\mathsf{E}_k])^{d-1}(H_{k+1}[N]G_k)$. Then as in case 1 it follows that $M \sim \mathsf{E}_k$. Moreover, by lemma 2.8 $H_{k+1}[\mathsf{E}_{2k}] =_{s<n} H_{k+1}[N]$, so by the induction hypothesis $\mathsf{E}_{2k} \sim N$. So $M \sim \mathsf{E}_k \sim \mathsf{E}_{2k} \sim N$.

Case 3. Both $W_M, W_N$ are reached after $\geq 8$ steps. Then

$$\begin{array}{lll} W_M & \equiv & D_k^j[\mathsf{E}_k]((H_{k+1}[\mathsf{E}_k])^d(H_{k+1}[M]G_k)); \\ W_N & \equiv & D_k^j[\mathsf{E}_k]((H_{k+1}[\mathsf{E}_k])^{d'}(H_{k+1}[N]G_k)). \end{array}$$

If $d = d'$, then by lemma 2.7

$$(H_{k+1}[\mathsf{E}_k])^d(H_{k+1}[M]G_k) =_{s<n} (H_{k+1}[\mathsf{E}_k])^d(H_{k+1}[N]G_k),$$

so

$$H_{k+1}[M] =_{s<n} H_{k+1}[N],$$

since $H_{k+1}[X]$ is always of order 0. Therefore by the induction hypothesis $M \sim N$.

If on the other hand, say, $d < d'$, then (writing $d' = d + e$)

$$\begin{array}{lll} W_M & \equiv & D_k^j[\mathsf{E}_k]((H_{k+1}[\mathsf{E}_k])^d(H_{k+1}[M] \qquad\qquad G_k \qquad\qquad )); \\ W_N & \equiv & D_k^j[\mathsf{E}_k]((H_{k+1}[\mathsf{E}_k])^d(H_{k+1}[\mathsf{E}_k]\ \underline{((H_{k+1}[\mathsf{E}_k])^{e-1}(H_{k+1}[N]G_k))}\,)). \end{array}$$

so

$$\begin{array}{rll} H_{k+1}[M] & =_{s<n} & H_{k+1}[\mathsf{E}_k] \\ G_k & =_{s<n} & (H_{k+1}[\mathsf{E}_k])^{e-1}(H_{k+1}[N]G_k), \end{array}$$

since $H_{k+1}[X]$ is always of order 0. Therefore by lemma 2.8

$$H_{k+1}[\mathsf{E}_{2k}] \;=_{s<n}\; H_{k+1}[N]$$

Therefore by the induction hypothesis twice we obtain $M \sim \mathsf{E}_k \sim \mathsf{E}_{2k} \sim N$. ■

## References

Barendregt, H. P. [1984]. *The Lambda Calculus: its syntax and semantics*, revised edition, North-Holland Publishing Co., Amsterdam.

Rogers, Hartley, Jr. [1987]. *Theory of recursive functions and effective computability*, second edition, MIT Press, Cambridge, Mass.

Statman, Rick [1986]. Every countable poset is embeddable in the poset of unsolvable terms, *Theoret. Comput. Sci.* **48**(1), pp. 95–100.

Statman, Rick [1998]. Morphisms and partitions of V-sets, *CSL'98*, Springer, Berlin. To appear.

Statman, Rick [1999]. Consequences of a theorem of Jacopini: consistent equalities and equations, *TLCA'99*, Springer, Berlin. To appear.

Visser, Albert [1980]. Numerations, $\lambda$-calculus & arithmetic, *To H. B. Curry: essays on combinatory logic, lambda calculus and formalism*, Academic Press, London, pp. 259–284.